

10/ 788 906

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number  
**WO 2006/019856 A1**

(51) International Patent Classification<sup>7</sup>: **A61N 1/30**

Minnesota 55125 (US). **ROSS, Jeffrey** [US/US]; 3106 Fairview Avenue, Roseville, MN 55113 (US).

(21) International Application Number:  
PCT/US2005/024914

(74) Agents: **STEFFEY, Charles E.** et al.; Schwegman, Lundberg, Woessner & Kluth, P.A., P.O. Box 2938, Minneapolis, Minnesota 55402 (US).

(22) International Filing Date: 13 July 2005 (13.07.2005)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
10/890,825 14 July 2004 (14.07.2004) US

(71) Applicant (for all designated States except US): **CARDIAC PACEMAKERS, INC.** [US/US]; 4100 Hamline Avenue North, St. Paul, Minnesota 55112-5798 (US).


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GIROUARD, Steven** [US/US]; 8155 Boulder Ridge Road, Woodbury,

[Continued on next page]

(54) Title: METHOD AND APPARATUS FOR CONTROLLED GENE OR PROTEIN DELIVERY



(57) Abstract: An implantable system which includes a gene/protein delivery device and a pulse generator, as well as method of preparing the gene/protein delivery device and using the system, are provided. In one embodiment, the implantable system detects a predetermined condition or event and, in response, delivers gene(s) and/or protein(s) in conjunction with delivering pacing and/or defibrillation pulses.

WO 2006/019856 A1



FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,  
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

— *with international search report*

## METHOD AND APPARATUS FOR CONTROLLED GENE OR PROTEIN DELIVERY

### Claim of Priority

5           Benefit of priority is hereby claimed to U.S. Patent Application Serial  
Number 10/890,825, filed on July 14, 2004, which application is herein  
incorporated by reference.

### Field of the Invention

10           This invention relates generally to gene therapy of living tissue and  
particularly, but not by way of limitation, to method and apparatus for delivering  
gene or protein using an implantable medical device.

### Background of the Invention

15           Many techniques currently exist for delivering drugs or other  
medicaments to body tissue. These include topical or transcutaneous  
administration where the drug is passively absorbed, or caused to pass, into or  
across the skin or other surface tissue; oral administration; injection directly into  
body tissue such as through an intramuscular injection or the like; and  
20   intravenous administration, which involves introducing a selected drug directly  
into the blood stream. Transcutaneous drug delivery systems are usually limited  
to external administration of a drug through the patient's skin or other surface  
tissue, and thus is inefficient because some of the drug may be absorbed by  
healthy tissue before it reaches the diseased or damaged area, or carried beyond  
25   the diseased or damaged area. Oral administration, injection, and intravenous  
administration are systemic and so fail to concentrate the drug in a local area.

          Transportation of a drug using a localized drug delivery system may be  
enhanced through means such as iontophoresis. Iontophoresis typically involves  
an interaction between ionized molecules of a drug and an external electric field,  
30   which results in the migration of charged molecules. The migration is  
accomplished by placing two electrodes across the tissue to be treated and  
charging the electrodes with a relatively low, direct current (DC), voltage. One  
of the electrodes acts as a source electrode and is typically in contact with the  
drug solution. The other electrode acts as a return electrode and may be filled

with an electrolyte solution. The electric field generated between the two electrodes causes the charged molecules to migrate from one electrode into the tissues to be treated.

Nevertheless, problems are associated with introducing an electrical  
5 current into the body, including muscle stimulation and contraction, as well as pain or other unwanted sensations. More importantly, the problem of cardiac arrhythmia (irregular rhythm) can easily arise when electrical current passes through the heart. The current source causing that problem can originate from an external source, within the heart itself, or adjacent to the heart, such as from a  
10 coronary artery.

Intensity (current density), frequency, waveform and duration of the electrical current used in iontophoresis have an effect on whether cardiac arrhythmias and other problems will occur, as well as the magnitude of those reactions. The threshold at which ventricular fibrillation occurs with various  
15 transthoracic and intracardiac electrical levels increases with higher frequency currents. The threshold of sensation also increases with higher frequencies. For instance, U.S. Patent No. 5,087,243 discloses a method which attempts to minimize the risk of iontophoresis-induced arrhythmias. An implanted myocardial iontophoresis patch system is disclosed therein which a pulsed  
20 current is supplied to the anodal patch. Electrical activity in the patient's heart is monitored and the iontophoresis current is pulsed on and off in synchronization with ventricular depolarization to avoid the interval during which the heart is vulnerable to electrically induced arrhythmias or unnatural heart rhythms.

What is needed is an improved apparatus for local, transient delivery of  
25 gene(s) and/or protein(s), e.g., one which is employed in conjunction with electrical therapy.

### **Summary of the Invention**

The present invention is directed to the local, transient delivery of a  
30 therapeutic agent, such as a nucleic acid molecule (polynucleotide) or protein, from a polymer matrix to the myocardium, a vessel, or any other organ or area for which transluminal access is desirable of a mammal having or at risk of a particular condition. For example, the present invention can be used to locally administer via iontophoresis isolated nucleic acid or protein to the myocardium

of a mammal having or at risk of a cardiovascular condition in response to detection of a parameter or a change in a parameter which is associated with the condition, e.g., the isolated nucleic acid and/or protein is administered in an amount effective to transiently alter, e.g., enhance, myocardial function.

- 5 Iontophoresis technology uses an electrical potential or current across a permeable barrier to drive ionic molecules such as nucleic acid or protein, or drag nonionic molecules in an ionic solution. Iontophoresis offers continuous or pulsatile delivery as well as preprogrammed administration schedules. Factors affecting iontophoretic transport include pH, current density, ionic strength,  
10 concentration of the molecule to be transported, molecular size of the molecule to be transported, and the method of current application (continuous or pulse current). The charge on the molecule can be controlled by changing the pH of the solution, and so delivery can be adjusted for either cathodal or anodal iontophoresis. Iontophoresis can thus facilitate transport of the molecule and  
15 may enhance tissue penetration.

In one embodiment, a mammal has a condition associated with aberrant expression, for instance, aberrant temporal expression or aberrant levels of expression, of a gene product, e.g., under expression or lack of expression of wild-type (functional) gene product. To prevent, inhibit or treat the condition in  
20 the mammal, an implantable device in the mammal releases via iontophoresis nucleic acid from a polymer matrix which includes at least one gene or a portion thereof in sense orientation, e.g., an open reading frame for the gene product or a portion thereof which encodes a gene product with substantially the same activity as the full-length gene product. In one embodiment, myocardial tissues  
25 of a mammal are contacted with genes which directly or indirectly modulate conduction in the myocardium, e.g., genes for gene products which modulate gap junction proteins and/or ion channel proteins, or for transcriptional regulatory proteins which alter gap junction protein or ion channel protein levels. In another embodiment, atria of mammal having an aberrant  $I_f$  are contacted with  
30 an implantable device which includes HCN nucleic acid operably linked to a promoter (an expression cassette) embedded in a polymer matrix. Upon detection of the aberrant  $I_f$ , the HCN gene is delivered to atrial cells in an amount which alters the  $I_f$ . Thus, for myocardial conditions, the isolated nucleic acid and/or protein embedded or applied to a polymer matrix is introduced to the

epicardium, endocardium, or pericardium, thereby providing for localized transmyocardial delivery at a lower dosage relative to systemic delivery. In another embodiment, the nucleic acid in the polymer matrix encodes a dominant negative gene product which is useful to decrease expression or levels of an endogenous molecule, for instance, one which is formed of multiple subunits, one of which subunits corresponds to the gene product encoded by the nucleic acid in the polymer matrix.

In another embodiment, an implantable device in a mammal having a condition associated with under expression of a wild-type (functional) protein releases via iontophoresis the protein from a polymer matrix in an amount which is effective to prevent, inhibit or treat the condition.

Antisense oligonucleotides or polynucleotides can inhibit gene expression, mainly by binding to messenger RNA of the target gene, and thus have the ability to block the expression of gene products associated with a particular condition. Thus, in one embodiment, the nucleic acid in the polymer matrix includes a gene or a portion thereof, e.g., an oligonucleotide, in antisense orientation. In one embodiment, the presence of the nucleic acid in a cell of a mammal with a corresponding endogenous gene inhibits the expression of the endogenous gene. In one embodiment, the endogenous gene encodes a wild-type (functional) gene product which is overexpressed. In another embodiment, the endogenous gene encodes a mutant gene product, e.g., one with altered activity or a dominant negative.

#### **Brief Description of the Drawings**

In the drawings, which are not necessarily drawn to scale, like numerals describe similar components throughout the several views. The drawings illustrate generally, by way of example, but not by way of limitation, various embodiments discussed in the present document. The drawings are for illustrative purposes only and not to scale nor anatomically accurate.

FIG. 1 is an illustration of an embodiment of a gene/protein delivery system and portions of an environment in which it is used.

FIG. 2 is a block diagram showing one embodiment of the circuit of portions of the gene/protein delivery system such as shown in FIG. 1.

FIG. 3 is a block diagram showing a further embodiment of the circuit of portions of the gene/protein delivery system such as shown in FIG. 1.

### **Detailed Description**

5 In the following detailed description, reference is made to the accompanying drawings which form a part hereof, and in which is shown by way of illustration specific embodiments in which the invention may be practiced. These embodiments are described in sufficient detail to enable those skilled in the art to practice the invention, and it is to be understood that the  
10 embodiments may be combined, or that other embodiments may be utilized and that structural, logical and electrical changes may be made without departing from the spirit and scope of the present invention. The following detailed description provides examples, and the scope of the present invention is defined by the appended claims and their equivalents.

15 It should be noted that references to "an", "one", or "various" embodiments in this disclosure are not necessarily to the same embodiment, and such references contemplate more than one embodiment.

### **Definitions**

By "nucleic acid", "oligonucleotide", and "polynucleotide" or  
20 grammatical equivalents herein means at least two nucleotides covalently linked together.

A "polynucleotide" refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides, or analogs thereof. This term refers to the primary structure of the molecule, and thus includes double-  
25 and single-stranded DNA, as well as double- and single-stranded RNA, and portions of both double stranded or single stranded sequence. The polynucleotide may be DNA, both genomic and cDNA, RNA or a hybrid, where the polynucleotide contains any combination of deoxyribo- and ribo-nucleotides, and any combination of bases, including uracil, adenine, thymine, cytosine,  
30 guanine, inosine, xanthine and hypoxanthine, etc. Thus, for example, chimeric DNA-RNA molecules may be used such as described in Cole-Strauss et al., Science, 273:1386 (1996) and Yoon et al., Proc. Natl. Acad. Sci. USA, 93:2071 (1996). It also includes modified polynucleotides such as methylated and/or capped polynucleotides.

An "oligonucleotide" includes at least 7 nucleotides, preferably 15, and more preferably 20 or more sequential nucleotides, up to 100 nucleotides, either RNA or DNA, which correspond to the complement of the non-coding strand, or of the coding strand, of a selected mRNA, or which hybridize to the mRNA or  
5 DNA encoding the mRNA and remain stably bound under moderately stringent or highly stringent conditions, as defined by methods well known to the art, e.g., in Sambrook et al., A Laboratory Manual, Cold Spring Harbor Press (1989).

The term "isolated" when used in relation to a nucleic acid, peptide or polypeptide refers to a nucleic acid sequence, peptide or polypeptide that is  
10 identified and separated from at least one contaminant nucleic acid, polypeptide or other biological component with which it is ordinarily associated in its natural source. Isolated nucleic acid, peptide or polypeptide is present in a form or setting that is different from that in which it is found in nature. For example, a given DNA sequence (e.g., a gene) is found on the host cell chromosome in  
15 proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs that encode a multitude of proteins. The isolated nucleic acid molecule may be present in single-stranded or double-stranded form. When an isolated nucleic acid molecule is to be utilized to express a protein, the  
20 molecule will contain at a minimum the sense or coding strand (i.e., the molecule may single-stranded), but may contain both the sense and anti-sense strands (i.e., the molecule may be double-stranded).

"Recombinant," as applied to a polynucleotide, means that the polynucleotide is the product of various combinations of cloning, restriction  
25 and/or ligation steps, and other procedures that result in a construct that is distinct from a polynucleotide found in nature. Recombinant as applied to a protein means that the protein is the product of expression of a recombinant polynucleotide.

A "gene" generally refers to a polynucleotide or portion of a  
30 polynucleotide which includes a sequence for a gene product. For most situations, it is desirable for the gene to also comprise a promoter operably linked to the coding sequence in order to effectively promote transcription. Enhancers, repressors and other regulatory sequences may also be included in order to modulate activity of the gene, as is well known in the art.



The terms "polypeptide," "peptide," and "protein" are used interchangeably to refer to polymers of amino acids of any length. These terms also include proteins that are post-translationally modified through reactions that include glycosylation, acetylation and phosphorylation.

5 By "DNA" is meant a polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in double-stranded or single-stranded form found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure of particular DNA molecules, sequences may be described herein according to the normal  
10 convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having the sequence complementary to the mRNA). The term captures molecules that include the four bases adenine, guanine, thymine, or cytosine, as well as molecules that include base analogues which are known in the art.

15 As used herein, the terms "complementary" or "complementarity" are used in reference to polynucleotides (i.e., a sequence of nucleotides) related by the base-pairing rules. For example, the sequence "A-G-T," is complementary to the sequence "T-C-A." Complementarity may be "partial," in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or,  
20 there may be "complete" or "total" complementarity between the nucleic acids. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, as well as detection methods that depend upon binding between nucleic acids.

25 DNA molecules are said to have "5' ends" and "3' ends" because mononucleotides are reacted to make oligonucleotides or polynucleotides in a manner such that the 5' phosphate of one mononucleotide pentose ring is attached to the 3' oxygen of its neighbor in one direction via a phosphodiester linkage. Therefore, an end of an oligonucleotide or polynucleotide is referred to  
30 as the "5' end" if its 5' phosphate is not linked to the 3' oxygen of a mononucleotide pentose ring and as the "3' end" if its 3' oxygen is not linked to a 5' phosphate of a subsequent mononucleotide pentose ring. As used herein, a nucleic acid sequence, even if internal to a larger oligonucleotide or polynucleotide, also may be said to have 5' and 3' ends. In either a linear or

circular DNA molecule, discrete elements are referred to as being "upstream" or 5' of the "downstream" or 3' elements. This terminology reflects the fact that transcription proceeds in a 5' to 3' fashion along the DNA strand. The promoter and enhancer elements that direct transcription of a linked gene are generally  
5 located 5' or upstream of the coding region. However, enhancer elements can exert their effect even when located 3' of the promoter element and the coding region. Transcription termination and polyadenylation signals are located 3' or downstream of the coding region.

"Homology" refers to the percent of identity between two  
10 polynucleotides or two polypeptides. The correspondence between one sequence and to another can be determined by techniques known in the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide or polynucleotide molecules by aligning the sequence information and using readily available computer programs. Alternatively,  
15 homology can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single strand-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide, sequences are "substantially homologous" to each other when at least about 80%, preferably at  
20 least about 90%, and most preferably at least about 95% of the nucleotides, or amino acids, respectively match over a defined length of the molecules, as determined using the methods above.

By "expression construct" or "expression cassette" is meant a nucleic acid molecule that is capable of directing transcription. An expression construct  
25 includes, at the least, a promoter. Additional elements, such as an enhancer, and/or a transcription termination signal, may also be included.

The term "heterologous" as it relates to nucleic acid sequences such as gene sequences and control sequences, denotes sequences that are not normally joined together, and/or are not normally associated with a particular cell. Thus, a  
30 "heterologous" region of a nucleic acid construct or a vector is a segment of nucleic acid within or attached to another nucleic acid molecule that is not found in association with the other molecule in nature. For example, a heterologous region of a nucleic acid construct could include a coding sequence flanked by sequences not found in association with the coding sequence in nature, i.e., a

heterologous promoter. Another example of a heterologous coding sequence is a construct where the coding sequence itself is not found in nature (e.g., synthetic sequences having codons different from the native gene). Similarly, a cell transformed with a construct which is not normally present in the cell would be considered heterologous for purposes of this invention.

The term "control elements" refers collectively to promoter regions, polyadenylation signals, transcription termination sequences, upstream regulatory domains, origins of replication, internal ribosome entry sites ("IRES"), enhancers, splice junctions, and the like, which collectively provide for the replication, transcription, post-transcriptional processing and translation of a coding sequence in a recipient cell. Not all of these control elements need always be present so long as the selected coding sequence is capable of being replicated, transcribed and translated in an appropriate host cell.

The term "promoter region" is used herein in its ordinary sense to refer to a nucleotide region comprising a DNA regulatory sequence, wherein the regulatory sequence is derived from a gene which is capable of binding RNA polymerase and initiating transcription of a downstream (3' direction) coding sequence. Thus, a "promoter," refers to a polynucleotide sequence that controls transcription of a gene or coding sequence to which it is operably linked. A large number of promoters, including constitutive, inducible and repressible promoters, from a variety of different sources, are well known in the art.

By "enhancer element" is meant a nucleic acid sequence that, when positioned proximate to a promoter, confers increased transcription activity relative to the transcription activity resulting from the promoter in the absence of the enhancer domain. Hence, an "enhancer" includes a polynucleotide sequence that enhances transcription of a gene or coding sequence to which it is operably linked. A large number of enhancers, from a variety of different sources are well known in the art. A number of polynucleotides which have promoter sequences (such as the commonly-used CMV promoter) also have enhancer sequences.

By "cardiac-specific enhancer or promoter" is meant an element, which, when operably linked to a promoter or alone, respectively, directs gene expression in a cardiac cell and does not direct gene expression in all tissues or all cell types. Cardiac-specific enhancers or promoters may be naturally occurring or non-naturally occurring. One skilled in the art will recognize that

the synthesis of non-naturally occurring enhancers or promoters can be performed using standard oligonucleotide synthesis techniques.

"Operably linked" refers to a juxtaposition, wherein the components so described are in a relationship permitting them to function in their intended manner. By "operably linked" with reference to nucleic acid molecules is meant that two or more nucleic acid molecules (e.g., a nucleic acid molecule to be transcribed, a promoter, and an enhancer element) are connected in such a way as to permit transcription of the nucleic acid molecule. A promoter is operably linked to a coding sequence if the promoter controls transcription of the coding sequence. Although an operably linked promoter is generally located upstream of the coding sequence, it is not necessarily contiguous with it. An enhancer is operably linked to a coding sequence if the enhancer increases transcription of the coding sequence. Operably linked enhancers can be located upstream, within or downstream of coding sequences. A polyadenylation sequence is operably linked to a coding sequence if it is located at the downstream end of the coding sequence such that transcription proceeds through the coding sequence into the polyadenylation sequence. "Operably linked" with reference to peptide and/or polypeptide molecules is meant that two or more peptide and/or polypeptide molecules are connected in such a way as to yield a single polypeptide chain, i.e., a fusion polypeptide, having at least one property of each peptide and/or polypeptide component of the fusion. Thus, a signal or targeting peptide sequence is operably linked to another protein if the resulting fusion is secreted from a cell as a result of the presence of a secretory signal peptide or into an organelle as a result of the presence of an organelle targeting peptide.

"Gene delivery," "gene transfer," and the like as used herein, are terms referring to the introduction of an exogenous polynucleotide (sometimes referred to as a "transgene") into a host cell, irrespective of the method used for the introduction. Such methods include a variety of well-known techniques such as vector-mediated gene transfer (by, e.g., viral infection/transfection, or various other protein-based or lipid-based gene delivery complexes) as well as techniques facilitating the delivery of "naked" polynucleotides (such as electroporation, iontophoresis, "gene gun" delivery and various other techniques used for the introduction of polynucleotides). The introduced polynucleotide may be stably or transiently maintained in the host cell. In one embodiment,

e.g., with isolated nucleic acid which is not extrachromosomally maintained or integrated into the genome of a cell, the introduced polynucleotide is transiently maintained in the cell. Stable maintenance typically requires that the introduced polynucleotide either contains an origin of replication compatible with the host cell or integrates into a replicon of the host cell such as an extrachromosomal replicon (e.g., a plasmid) or a nuclear or mitochondrial chromosome.

By "transgene" is meant any piece of a nucleic acid molecule (for example, DNA) which is inserted by artifice into a cell either transiently or permanently, and becomes part of the organism if integrated into the genome or maintained extrachromosomally. Such a transgene may include a gene which is partly or entirely heterologous (i.e., foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

"*In vivo*" gene/protein delivery, gene/protein transfer, gene/protein therapy and the like as used herein, are terms referring to the introduction of an exogenous (isolated) polynucleotide or protein directly into the body of an organism, such as a human or non-human mammal, whereby the exogenous polynucleotide or protein is introduced to a cell of such organism *in vivo*.

A "vector" (sometimes referred to as gene delivery or gene transfer "vehicle") refers to a macromolecule or complex of molecules comprising a polynucleotide or protein to be delivered to a host cell, either *in vitro* or *in vivo*. Vectors include, for example, expression vectors, viral vectors (such as adenoviruses ("Ad"), adeno-associated viruses (AAV), lentiviruses, poxviruses, papilloma viruses, herpesviruses, and retroviruses), liposomes and other lipid-containing complexes, and other macromolecular complexes capable of mediating or enhancing delivery of a polynucleotide or protein to a host cell. Vectors can also comprise other components or functionalities that further modulate gene delivery and/or gene expression, or that otherwise provide beneficial properties to the targeted cells. Such other components include, for example, components that influence binding or targeting to cells (including components that mediate cell-type or tissue-specific binding); components that influence uptake of nucleic acid or protein by the cell; components that influence localization of the polynucleotide or protein within the cell after uptake (such as agents mediating nuclear localization); and components that influence expression of polynucleotides. Such components also might include markers,

such as detectable and/or selectable markers that can be used to detect or select for cells that have taken up and are expressing the introduced nucleic acid. Such components can be provided as a natural feature of the vector (such as the use of certain viral vectors which have components or functionalities mediating binding and uptake), or vectors can be modified to provide such functionalities. A large variety of such vectors are known in the art and are generally available.

A "therapeutic polynucleotide", "therapeutic gene" or "therapeutic protein" refers to a nucleotide sequence or amino acid sequence that is capable, when transferred to cells of an individual, of eliciting a prophylactic, curative or other beneficial effect in the individual.

The term "corresponds to" is used herein to mean that a polynucleotide or protein sequence is homologous (i.e., may be similar or identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide or protein sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary polynucleotide sequence is able to hybridize to the other strand. As outlined below, preferably, the homology between the two sequences is at least 70%, preferably 85%, and more preferably 95%, identical.

The terms "substantially corresponds to" or "substantial identity" or "homologous" as used herein denotes a characteristic of a nucleic acid or protein sequence, wherein a nucleic acid or protein sequence has at least about 70% sequence identity as compared to a reference sequence, typically at least about 85% sequence identity, and preferably at least about 95% sequence identity, as compared to a reference sequence. The reference sequence may be a subset of a larger sequence, such as a portion of a gene or flanking sequence, or portion of protein. However, the reference sequence is at least 20 nucleotides long, typically at least about 30 nucleotides long, and preferably at least about 50 to 100 nucleotides long, or, for peptides or polypeptides, at least 7 amino acids long, typically at least 10 amino acids long, and preferably at least 20 to 30 amino acids long. "Substantially complementary" as used herein refers to a nucleotide sequence that is complementary to a sequence that substantially corresponds to a reference sequence.

"Specific hybridization" is defined herein as the formation of hybrids between a polynucleotide which may include substitutions, deletion, and/or additions as compared to a reference sequence and a selected target nucleic acid

sequence, wherein the polynucleotide preferentially hybridizes to a target nucleic acid sequence such that, for example, at least one discrete band can be identified on a Northern or Southern blot of DNA prepared from cells that contain the target nucleic acid sequence. It is evident that optimal hybridization conditions will vary depending upon the sequence composition and length(s) of the polynucleotide(s) and target(s), and the experimental method selected by the practitioner. Various guidelines may be used to select appropriate hybridization conditions (see, Maniatis et al., 1989 and Berger and Kimmel, 1987).

The term "wild-type" refers to a gene or gene product that has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designated the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product that displays modifications in sequence and or functional properties (i.e., altered characteristics) when compared to the wild-type gene or gene product. It is noted that naturally-occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "disease allele" refers to an allele of a gene that is capable of producing a recognizable disease. A disease allele may be dominant or recessive and may produce disease directly or when present in combination with a specific genetic background or pre-existing pathological condition. A disease allele may be present in the gene pool (an inherited disease allele) or may be generated *de novo* in an individual by somatic mutation (an acquired disease allele).

"Vasculature" or "vascular" are terms referring to the system of vessels carrying blood (as well as lymph fluids) throughout the mammalian body.

"Blood vessel" refers to any of the vessels of the mammalian vascular system, including arteries, arterioles, capillaries, venules, veins, sinuses, and vasa vasorum.

"Artery" refers to a blood vessel through which blood passes away from the heart. Coronary arteries supply the tissues of the heart itself, while other arteries supply the remaining organs of the body. The general structure of an artery consists of a lumen surrounded by a multi-layered arterial wall.

By "mammal" is meant any member of the class *Mammalia* including, without limitation, humans and nonhuman primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals  
5 including rodents such as mice, rats, rabbits and guinea pigs, and the like.

By "derived from" is meant that a nucleic acid molecule was either made or designed from a parent nucleic acid molecule, the derivative retaining substantially the same functional features of the parent nucleic acid molecule, e.g., encoding a gene product with substantially the same activity as the gene  
10 product encoded by the parent nucleic acid molecule from which it was made or designed.

The term "exogenous," when used in relation to a protein, gene, nucleic acid, or polynucleotide in a cell or organism refers to a protein, gene, nucleic acid, or polynucleotide which has been introduced into the cell or organism by  
15 artificial or natural means, or in relation a cell refers to a cell which was isolated and subsequently introduced to other cells or to an organism by artificial or natural means. An exogenous nucleic acid may be from a different organism or cell, or it may be one or more additional copies of a nucleic acid which occurs naturally within the organism or cell. An exogenous cell may be from a different  
20 organism, or it may be from the same organism. By way of a non-limiting example, an exogenous nucleic acid is in a chromosomal location different from that of natural cells, or is otherwise flanked by a different nucleic acid sequence than that found in nature.

By "growth factor" is meant an agent that, at least, promotes cell growth  
25 or induces phenotypic changes.

The term "angiogenic" means an agent that alone or in combination with other agents induces angiogenesis, and includes, but is not limited to, fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), hepatocyte growth factor, angiogenin, transforming growth factor (TGF), tissue necrosis  
30 factor (TNF, e.g., TNF- $\alpha$ ), platelet derived growth factor (PDGF), granulocyte colony stimulatory factor (GCSF), placental GF, IL-8, proliferin, angiopoietin, e.g., angiopoietin-1 and angiopoietin-2, thrombospondin, ephrin-A1, E-selectin, leptin and heparin affinity regulatory peptide.



“Cardiovascular conditions” include, but are not limited to, coronary artery disease/ischemia, coronary artery disease (CAD), ischemia, angina (chest pain), thrombosis, coronary thrombosis, myocardial infarction (MI), silent ischemia, stenosis/restenosis, transient ischemic attack (TIA), atherosclerosis,

5 peripheral vascular disease, bradyarrhythmia, e.g., bradyarrhythmia, bradycardia, sick sinus rhythm (Sick Sinus Syndrome), sinus bradycardia, sinoatrial block, asystole, sinus arrest, syncope, first degree atrio-ventricular (AV) block, second degree atrio-ventricular (AV) block, third degree atrio-ventricular (AV) block,

10 chronotropic incompetence, tachyarrhythmia, e.g., tachyarrhythmia, tachycardia, fibrillation, flutter, atrial fibrillation, atrial flutter, familial atrial fibrillation, paroxysmal atrial fibrillation, permanent atrial fibrillation, persistent atrial fibrillation, supraventricular tachyarrhythmias, sinus tachycardia, reentry (reentrant arrhythmias), AV nodal reentry, focal arrhythmia, ectopy, ventricular

15 fibrillation (VF), ventricular tachycardia (VT), Wolff-Parkinson-White Syndrome (WPW) and sudden cardiac death, heart failure, e.g., heart failure, cardiomyopathy, congestive heart failure, hypertrophic cardiomyopathy, remodeling, non-ischemic cardiomyopathy, dilated cardiomyopathy, restrictive

20 cardiomyopathy, diastolic heart failure, systolic heart failure, and chronic heart failure, heart block/electrical disorders, e.g., atrioventricular (AV) block, bundle branch block (BBB), left bundle branch block (LBBB), right bundle branch

block (RBBB), Long QT Syndrome (LQTS), premature ventricular contraction (PVC), electrical remodeling, intraventricular conduction defect, and hemiblock, hemodynamic deficiency, e.g., hypertension, hypotension, left ventricular

25 dysfunction, low ejection fraction, low cardiac output, and low stroke volume, sudden cardiac death, cardiac arrest, sudden cardiac death (SCD), ventricular fibrillation, and pump failure, as well as bacterial endocarditis, viral myocarditis, pericarditis, rheumatic heart disease, and syncope. In particular, a

cardiovascular condition includes, but is not limited to, arrhythmia, e.g., atrial fibrillation, ventricular fibrillation or bradycardia, ischemia, heart failure and

30 hyperplasia not associated with neoplastic disease, which condition may be associated with ventricular remodeling, diastolic dysfunction, aberrant body temperature, aberrant or altered pressure, e.g., altered venous, left ventricular or left atrial pressure, aberrant or altered heart rate or sounds, aberrant or altered

electrogram, aberrant or altered cardiac metabolism, such as altered blood pH,

glucose, pO<sub>2</sub>, pCO<sub>2</sub>, minute ventilation, creatine, CRP, Mef2A, creatine kinase or creatine kinase MB levels, aberrant or altered pulmonary or thoracic impedance, aberrant or altered stroke volume, aberrant or altered neurohormone levels, aberrant or altered electrical activity, aberrant or altered sympathetic  
5 nerve activity, aberrant or altered renal output, aberrant or altered filtration rate, aberrant or altered angiotensin II levels, or aberrant or altered respiratory sounds, and the like.

"Treatment" or "therapy" as used herein refers to administering, to an individual patient, agents that are capable of eliciting a prophylactic, curative or  
10 other beneficial effect in the individual.

"Gene therapy" as used herein refers to administering, to an individual patient, vectors comprising a therapeutic gene.

A "user" includes a physician or other caregiver using the gene/protein delivery system to treat a patient.

15

#### General Overview

The present invention provides a system to transiently deliver one or more gene- and/or one or more protein-based therapeutic agents, as well as nongene-, nonprotein-based therapeutic agents, in a spatially controlled manner  
20 to cells or tissue of a mammal. The isolated nucleic acid, when introduced and optionally expressed in a cell, or the isolated protein when introduced to a cell, yields a cell with a different phenotype than a corresponding cell which is not contacted with the isolated nucleic acid or isolated protein. In some embodiments, the isolated nucleic acid or protein will be other than a naturally  
25 occurring sequence. In one embodiment, the mammal to be treated has or is at risk of having a cardiovascular condition. For example, the mammal may have or be at risk of having atrial fibrillation, ventricular tachycardia, heart failure, ischemia, bradycardia or hyperplasia.

This document also discusses a gene/protein delivery system that  
30 includes an implantable iontophoresis gene/protein delivery device and an implantable pulse generator. In this document, "gene/protein delivery" includes delivery of genes, proteins, and optionally other moieties useful to inhibit, prevent or treat at least one symptom (manifestation) of a condition, e.g., a cardiovascular condition. The implantable iontophoresis gene/protein delivery

device is capable of delivering one or more genes and/or one or more proteins and optionally one or more non-nucleic acid, non-protein based agents to a mammal. As used herein, a "gene" includes a polynucleotide which may include a full-length open reading frame which encodes a gene product (sense orientation) or a portion thereof (sense orientation) which encodes a gene product with substantially the same activity as the gene product encoded by the full-length open reading frame, the complement of the polynucleotide, e.g., the complement of the full-length open reading frame (antisense orientation) and optionally linked 5' and/or 3' noncoding sequence(s) or a portion thereof, e.g., an oligonucleotide, which is useful to inhibit transcription, stability or translation of a corresponding mRNA. In one embodiment, a polynucleotide of the invention includes operably linked transcriptional elements including, but not limited to, a promoter, an enhancer, an intron, a Kozak sequence, or a transcription termination sequence. In another embodiment, a polynucleotide of the invention does not include a transcriptional element. In one embodiment, a polynucleotide of the invention includes modified nucleotides, e.g., those with one or more modified sugars, bases or phosphate groups. Thus, a nucleic acid of the present invention may contain phosphodiester bonds, although in some cases nucleic acid analogs are included that may have alternate backbones, comprising, for example, phosphoramidate (Beaucage et al., Methods Mol. Biol., 20:33 (1993); Letsinger, J. Org. Chem., 35:3800 (1970); Sprinzl et al., Eur. J. Biochem., 81:579 (1977); Letsinger et al., Nucl. Acids Res., 14:3487 (1986)), phosphorothioate, phosphorodithioate, O-methylphosphoroamidite linkages, including morpholino modified oligo- or poly-nucleotides (Kang et al., Biopolymers, 32:1351 (1992); Summerton et al., Antisense Nucl. Acid. Drug Dev., 7:187 (1997); U.S. Patent Nos. 5,142,047 and 5,185,444), and peptide nucleic acid backbones and linkages (Nielsen, Orig. Life Evol. Biosph., 23:323 (1993)). Modifications of the ribose-phosphate backbone or bases may be done to facilitate the addition of other moieties such as chemical constituents, including 2' O-methyl and 5' modified substituents, and/or to increase the stability and half-life of such molecules in physiological environments. In particular, for antisense nucleic acid which is not expressed from a vector, base modifications (Herdewijn et al., Antisense Nucl. Acid. Drug Dev., 10:297 (2000); Mangos et al., Curr. Topics Med. Chem., 7, 1147 (2002)); sugar

modifications (Urban et al., Farmaco, 58:243 (2003)), e.g., arabinose derivatives (Wilds et al., Bioconj. Chem., 10:299 (1999)); and phosphate modifications such as 2'-methoxyethoxy (Kimber et al., Reprod. Biomed. Online, 6:318 (2003)); and/or the use of 5-(N-aminohexyl)carbamoyl-2'-O-methyluridine (Ito et al., 5 NAR, 31:2514 (2003)), may be employed to enhance stability *in vivo*. Moreover, the nucleic acid may be a chimera, e.g., made up of DNA and RNA, e.g., deoxyribonucleotides and ribonucleotides, e.g., 2'-O-methyl-ribonucleotides (see published U.S. patent application 20030045830).

The isolated nucleic acid or protein may be prepared via recombinant 10 means or chemical synthesis. For example, the isolated nucleic acid may be a plasmid, viral vector, cosmid, phage, BAC, YAC or other artificial chromosome, isolated nucleic acid from a cell transfected with a plasmid or other vector or infected with a recombinant virus, or *in vitro* transcribed nucleic acid. As described above, in one embodiment, the isolated nucleic acid includes modified 15 nucleotides, e.g., those which are more stable to hydrolysis, which nucleic acid is prepared via chemical synthesis. In one embodiment, the isolated nucleic acid is capable of being transcribed by RNA polymerase and hybridizing to target nucleic acid while in other embodiments, the isolated nucleic acid is capable of hybridizing to target nucleic acid, e.g., antisense nucleic acid, but not of being 20 transcribed *in vivo*.

A protein useful in the system and methods of the invention can be synthesized *in vitro*, e.g., by the solid phase peptide synthetic method or by recombinant DNA approaches. The solid phase peptide synthetic method is an established and widely used method, which is described in the following 25 references: Stewart et al., Solid Phase Peptide Synthesis, W. H. Freeman Co., San Francisco (1969); Merrifield, J. Am. Chem. Soc., 85 2149 (1963); Meienhofer in "Hormonal Proteins and Peptides," ed.; C.H. Li, Vol. 2 (Academic Press, 1973), pp. 48-267; Bavaay and Merrifield, "The Peptides," eds. E. Gross and F. Meienhofer, Vol. 2 (Academic Press, 1980) pp. 3-285; and 30 Clark-Lewis et al., Meth. Enzymol., 287, 233 (1997). These proteins can be further purified by fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on an anion-exchange resin such as DEAE; chromatofocusing; SDS-PAGE;

ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; or ligand affinity chromatography.

Once isolated and characterized, derivatives, e.g., chemically derived derivatives, of a given protein can be readily prepared. For example, amides of proteins may also be prepared by techniques well known in the art for converting a carboxylic acid group or precursor, to an amide. A preferred method for amide formation at the C-terminal carboxyl group is to cleave the protein from a solid support with an appropriate amine, or to cleave in the presence of an alcohol, yielding an ester, followed by aminolysis with the desired amine.

Salts of carboxyl groups of a protein may be prepared in the usual manner by contacting the protein with one or more equivalents of a desired base such as, for example, a metallic hydroxide base, e.g., sodium hydroxide; a metal carbonate or bicarbonate base such as, for example, sodium carbonate or sodium bicarbonate; or an amine base such as, for example, triethylamine, triethanolamine, and the like.

N-acyl derivatives of an amino group of a protein may be prepared by utilizing an N-acyl protected amino acid for the final condensation, or by acylating a protected or unprotected protein. O-acyl derivatives may be prepared, for example, by acylation of a free hydroxy protein or protein resin. Either acylation may be carried out using standard acylating reagents such as acyl halides, anhydrides, acyl imidazoles, and the like. Both N- and O-acylation may be carried out together, if desired.

Formyl-methionine, pyroglutamine and trimethyl-alanine may be substituted at the N-terminal residue of the protein. Other amino-terminal modifications include aminooxypentane modifications (see Simmons et al., *Science*, 276, 276 (1997)).

In addition, the amino acid sequence of a protein can be modified so as to result in a variant. The modification includes the substitution of at least one amino acid residue in the protein for another amino acid residue, including substitutions which utilize the D rather than L form, as well as other well known amino acid analogs, e.g., unnatural amino acids such as  $\alpha$ ,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, and the like. These analogs include phosphoserine, phosphothreonine, phosphotyrosine, hydroxyproline, gamma-carboxyglutamate; hippuric acid, octahydroindole-2-carboxylic acid, statine,

1,2,3,4,-tetrahydroisoquinoline-3-carboxylic acid, penicillamine, ornithine, citrulline,  $\alpha$ -methyl-alanine, para-benzoyl-phenylalanine, phenylglycine, propargylglycine, sarcosine,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\omega$ -N-methylarginine, and other similar amino acids and imino acids and tert-butylglycine.

One or more of the residues of the protein can be altered, so long as the resulting variant has substantially the same activity as that of the unmodified (functionally active) protein. For example, it is preferred that the variant has at least about 80% or more, at least 90%, the biological activity of the corresponding unmodified protein. Conservative amino acid substitutions are preferred. For example, aspartic-glutamic as acidic amino acids; lysine/arginine/histidine as basic amino acids; leucine/isoleucine/methionine/alanine/valine/glycine as hydrophobic amino acids; serine/threonine as hydrophilic amino acids. Conservative amino acid substitution also includes groupings based on side chains. In another example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. For example, it is reasonable to expect that replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the properties of the resulting variant. Whether an amino acid change results in a functional protein can readily be determined by assaying the specific activity of the variant.

Amino acid substitutions falling within the scope of the invention, are, in general, accomplished by selecting substitutions that do not differ significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, (b) the charge or hydrophobicity of the molecule at the target

site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- 5 (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic; trp, tyr, phe.

The invention also envisions variants with non-conservative  
10 substitutions. Non-conservative substitutions entail exchanging a member of one of the classes described above for another.

Acid addition salts of a protein or of amino residues of the protein may be prepared by contacting the protein or amine with one or more equivalents of the desired inorganic or organic acid, such as, for example, hydrochloric acid.

15 Esters of carboxyl groups of the proteins may also be prepared by any of the usual methods known in the art.

Moreover, it is also envisioned that the proteins are modified in a manner that increases their stability *in vivo*, e.g., their half-life or bioavailability.

Methods to prepare such derivatives are well known to the art. One method to  
20 stabilize peptides is to prepare derivatives which are cyclized peptides (see EPA 471,453 (amide bonds), such as that between lysine and aspartic acid side chains; EPA 467,701 (disulfide bonds); EPA 467,699 (thioether bonds). Other modifications which may increase *in vivo* stability are disclosed in Jameson et al. (Nature, 368, 744 (1994)); U.S. Patent No. 4,992,463; U.S. Patent No. 5,596,078  
25 and U.S. Patent No. 5,091,396.

The gene/protein delivery system of the invention detects a certain physiological signal, e.g., a change in a particular physiological parameter, indicative of a condition or the onset of a condition, e.g., a condition treatable by gene and/or protein therapy, and, in response to the detection, delivers an  
30 effective amount of the gene(s) and/or protein(s). Though discussed specifically as part of a cardiac rhythm management system, the gene/protein delivery system may be employed for all *in vivo* gene/protein therapies.

FIG. 1 is an illustration of an embodiment of a gene/protein delivery system 100 and portions of an environment in which it is used. System 100

includes implantable system 105, external system 155, and telemetry link 140 providing for communication between implantable system 105 and external system 155.

Implantable system 105 includes, among other things, implantable CRM device 110, lead system 108, and implantable iontophoresis gene/protein delivery device 130. As shown in FIG. 1, implantable CRM device 110 is implanted in body 102. In one embodiment, implantable CRM device 110 includes a gene/protein delivery controller. Gene/protein delivery device 130 is attached to heart 101. Lead system 108 includes one or more leads providing for communication between implantable CRM device 110 and gene/protein delivery device 130. In one embodiment, lead system 108 provides for wired communication between implantable CRM device 110 and gene/protein delivery device 130. In one embodiment, lead system 108 provides for communication between implantable CRM device 110 and gene/protein delivery device 130 through tissue conduction of an electric signal. In various embodiments, implantable CRM device 110 also includes a pacemaker, a cardioverter/defibrillator, a cardiac resynchronization therapy (CRT) device, a cardiac remodeling control therapy (RCT) device, a drug delivery device or a drug delivery controller, a cell therapy device, or any other implantable medical device. Lead system 108 further includes leads for sensing physiological signals and delivering pacing pulses, cardioversion/defibrillation shocks, and/or pharmaceutical or other substances.

External system 155 includes external device 150, network 160, and remote device 170. External device 150 is within the vicinity of implantable CRM device 110 and communicates with implantable CRM device 110 bi-directionally via telemetry link 140. Remote device 170 is in a remote location and communicates with external device 150 bi-directionally via network 160, thus allowing a user to monitor and treat a patient from a distant location.

System 100 allows gene/protein delivery to be triggered by any one of implantable CRM device 110, external device 150, or remote device 170. In one embodiment, implantable CRM device 110 triggers gene/protein delivery upon detecting a predetermined signal or condition. In another embodiment, external device 150 or remote device 170 triggers gene/protein delivery upon detecting a condition from a signal transmitted from implantable CRM device 110. In one



specific embodiment, external system 155 includes a processor running a therapy decision algorithm to determine whether and when to trigger gene/protein delivery. In another specific embodiment, external system 155 includes a user interface to present signals acquired by implantable CRM device 155 and/or the  
5 detected abnormal condition to a user and receives commands from the user for triggering gene/protein delivery. In another specific embodiment, the user interface includes a user input incorporated into external device 150 to receive commands from the user and/or the patient treated with system 100. For example, the patient may be instructed to enter a command for a gene/protein  
10 delivery when he senses certain symptoms, and another person near the patient may do the same upon observing the symptoms.

FIG. 2 is a block diagram showing one embodiment of the circuit of portions of system 100 including implantable CRM device 110, lead system 108, and gene/protein delivery device 130. In one embodiment, lead system 108  
15 provides for an electrical connection between implantable CRM device 110 and gene/protein delivery device 130, such that the implantable CRM device transmits a voltage or current signal to the gene/protein delivery device to control gene/protein delivery. In another embodiment, lead system 108 allows an electrical field to be created at or near gene/protein delivery device 130 by an  
20 electrical signal generated from implantable CRM device 110 to control gene/protein delivery...

Gene/protein delivery device 130 includes polymer matrix 232 and isolated nucleic acid which encodes at least one gene product, isolated nucleic acid which binds at least one selected mRNA, or isolated protein, or a  
25 combination thereof. The isolated nucleic acid and/or protein are released from gene/protein delivery device 232 in response to the gene/protein delivery control signal transmitted from implantable CRM device 110. In one embodiment, gene/protein delivery device 130 includes an epicardial patch for delivering the gene(s)/protein(s) into cardiac tissue. In another embodiment, gene/protein  
30 delivery device 130 is incorporated into a vascular device such as a stent to release the gene(s) and/or protein(s) into the blood. Examples of a drug delivery device that employs iontophoresis are described in U.S. Patent No. 5,041,107, "ELECTRICALLY CONTROLLABLE NON-OCCLUDING, BODY IMPLANTABLE DRUG DELIVERY SYSTEM" and U.S. Patent No.

6,689,117, "DRUG DELIVERY SYSTEM FOR IMPLANTABLE MEDICAL DEVICE", both assigned to Cardiac Pacemakers, Inc., which are incorporated herein by reference in their entirety.

Implantable CRM device 110 includes sensor 212, event detector 213,  
5 and implant controller 214. Sensor 212 senses a physiological signal indicative of a condition which may be inhibited or treated with gene/protein delivery. Event detector 213 detects the condition. Implant controller 214 includes gene/protein delivery control module 215, which transmits a signal to gene/protein delivery device 130 to trigger gene/protein delivery in response to a  
10 detected condition. In one embodiment, gene/protein delivery device 130 includes polymer matrix 232 that contains one or more genes, e.g., two or more different, and/or one or more proteins, and releases the gene(s) and/or protein(s) in response to an applied electrical field. In one embodiment, an electrical field is applied to polymer matrix 232 as a voltage delivered through lead system 108.  
15 In another embodiment, such as when gene/protein delivery device 130 is not directly wired to an implantable CRM device, an electrical field is applied to polymer matrix 232 by conduction through the tissue of body 102.

In one embodiment, sensor 212 includes a cardiac sensing circuit that senses one or more electrograms, and event detector 213 detects an arrhythmia  
20 from the one or more electrograms. In one embodiment, event detector 213 detects the arrhythmia by detecting heart rate and comparing the heart rate to one or more threshold rates. A bradycardia condition is detected when the heart rate falls below a bradycardia threshold. A tachycardia condition is detected when the heart rate exceeds a tachycardia threshold. In a further embodiment, event  
25 detector 213 detects the arrhythmia by comparing morphological features of the electrogram to one or more predetermined templates. Event detector 213 includes one or more of a bradycardia detector, tachycardia detector, fibrillation detector, and any other arrhythmia detectors. In one specific embodiment, event detector 213 includes an atrial fibrillation detector. In another specific  
30 embodiment, event detector 213 includes a ventricular fibrillation detector.

In one embodiment, sensor 212 senses a physiological signal indicative of ischemia, and event detector 213 includes an ischemia detector. In one specific embodiment, sensor 212 senses an electrogram and event detector 213 runs an automatic ischemia detection algorithm to detect an ischemic condition

from the electrogram. One specific example of an electrogram-based ischemia detector is discussed in Zhu et al., U.S. Patent Application No. 09/962,852, "EVOKED RESPONSE SENSING FOR ISCHEMIA DETECTION," filed on September 25, 2001, assigned to Cardiac Pacemakers, Inc., which is

5 incorporated herein by reference in its entirety. In another embodiment, sensor 212 includes an electrical impedance based sensor using a low carrier frequency (e.g., 100 Hz), and event detector 213 runs an automatic ischemia detection algorithm to detect an ischemic condition from the electrical impedance signal. Tissue electrical impedance has been shown to increase significantly during

10 ischemia, as discussed in Min et al. (International Journal of Bioelectromagnetism, 5:53 (2003)). Sensor 212 senses low frequency electrical impedance signal between electrodes interposed in the heart. Event detector 213 detects the ischemia as abrupt changes in impedance (such as abrupt increases in value). In another specific embodiment, sensor 212 includes a local heart

15 motion based sensor utilizing an accelerometer located within a lead body positioned on or in the heart, and event detector 213 runs an automatic ischemia detection algorithm to detect an ischemic condition from the acceleration signal. Event detector 213 detects ischemia as an abrupt decrease in the amplitude of local cardiac accelerations.

20 In one embodiment, sensor 212 includes a metabolic sensor that senses a metabolic signal indicative of a cardiac metabolic level (rate of metabolism of cardiac cells). Examples of the metabolic sensor include but are not limited to a pH sensor, an oxygen pressure (PO<sub>2</sub>) sensor, a carbon dioxide pressure (PCO<sub>2</sub>) sensor, a glucose sensor, a creatine sensor, a C-creative protein sensor, a creatine

25 kinase sensor, a creatine kinase-MB sensor, or any combination of such sensors. Event detector 213 determines the cardiac metabolic level from the metabolic signal and compares the cardiac metabolic level to one or more predetermined thresholds defining a normal range of cardiac metabolic level. An abnormal condition is detected when the cardiac metabolic level is outside of the normal

30 range of cardiac metabolic level.

In one embodiment, sensor 212 includes an implantable impedance sensor to measure pulmonary impedance, or impedance of a portion of the thoracic cavity. Event detector 213 detects an abnormal condition when the impedance is out of its normal range. For example, pulmonary edema, i.e., fluid

retention in the lungs resulting from the decreased cardiac output, increases pulmonary or thoracic impedance. In one specific embodiment, event detector 213 produces a signal when the pulmonary or thoracic impedance exceeds a predetermined threshold impedance. In one embodiment, the impedance sensor  
5 is a respiratory sensor that senses the patient's minute ventilation. An example of an impedance sensor sensing minute ventilation is discussed in U.S. Patent No. 6,459,929, "IMPLANTABLE CARDIAC RHYTHM MANAGEMENT DEVICE FOR ASSESSING STATUS OF CHF PATIENTS," assigned to Cardiac Pacemakers, Inc., which is incorporated herein by reference in its  
10 entirety.

In one embodiment, sensor 212 includes a pressure sensor. Abnormal conditions including arrhythmias and heart failure cause pressures in various portions of the cardiovascular system to deviate from their normal ranges. Event detector 213 detects the abnormal condition when a pressure is outside of its  
15 normal range. In one specific embodiment, event detector 213 includes a systolic dysfunction detector to detect an abnormal condition related to pressure during the systolic phase of a cardiac cycle. In another specific embodiment, event detector 213 includes a diastolic dysfunction detector to detect an abnormal condition related to pressure during the diastolic phase of a cardiac  
20 cycle. Examples of the pressure sensor include but are not limited to a left atrial (LA) pressure sensor, a left ventricular (LV) pressure sensor, an artery pressure sensor, and a pulmonary artery pressure sensor. Pulmonary edema results in elevated LA and pulmonary arterial pressures. A deteriorated LV results in decreased LV and arterial pressures. In various embodiments, event detector  
25 213 detects an abnormal condition when the LA pressure exceeds a predetermined threshold LA pressure level, when the pulmonary arterial pressure exceeds a predetermined threshold pulmonary arterial pressure level, when the LV pressure falls below a predetermined threshold LV pressure level, and/or when the arterial pressure falls below a predetermined threshold LV pressure  
30 level. In other embodiments, event detector 213 derives a parameter from one of these pressures, such as a rate of change of a pressure, and produces a signal when the parameter deviates from its normal range. In one embodiment, the LV pressure sensor senses the LV pressure indirectly, by sensing a signal having known or predictable relationships with the LV pressure during all or a portion

of the cardiac cycle. Examples of such a signal include but are not limited to an LA pressure and a coronary vein pressure. One specific example of measuring the LV pressure using a coronary vein pressure sensor is discussed in U.S. Patent Application Serial No. 10/038,936, "METHOD AND APPARATUS FOR  
5 MEASURING LEFT VENTRICULAR PRESSURE," filed on January 4, 2002, assigned to Cardiac Pacemakers, Inc., which is hereby incorporated by reference in its entirety.

In one embodiment, sensor 212 includes a cardiac output or stroke volume sensor. Examples of stroke volume sensing are discussed in U.S. Patent  
10 No. 4,686,987, "BIOMEDICAL METHOD AND APPARATUS FOR CONTROLLING THE ADMINISTRATION OF THERAPY TO A PATIENT IN RESPONSE TO CHANGES IN PHYSIOLOGIC DEMAND," and U.S. Patent No. 5,284,136, "DUAL INDIFFERENT ELECTRODE PACEMAKER," both assigned to Cardiac Pacemakers, Inc., which are incorporated herein by  
15 reference in their entirety. Event detector 213 detects the abnormal condition when the stroke volume falls below a predetermined threshold level.

In one embodiment, sensor 212 includes a neural activity sensor to detect activities of the sympathetic nerve and/or the parasympathetic nerve. A significant decrease in cardiac output immediately stimulates sympathetic and/or  
20 the parasympathetic activities, as the autonomic nervous system attempts to compensate for deteriorated cardiac function. In one specific embodiment, a neural activity sensor includes a neurohormone sensor to sense a neurohormone level. Event detector 213 detects the abnormal condition when the hormone level exceeds a predetermined threshold level. In another specific embodiment,  
25 a neural activity sensor includes an action potential recorder to sense the electrical activities in the sympathetic nerve and/or the parasympathetic nerve. Event detector 213 detects an abnormal condition when the frequency of the electrical activities in the sympathetic and/or the parasympathetic nerves exceed a predetermined threshold level. Examples of direct and indirect neural activity  
30 sensing are discussed in U.S. Patent No. 5,042,497, "ARRHYTHMIA PREDICTION AND PREVENTION FOR IMPLANTED DEVICES," assigned to Cardiac Pacemakers, Inc., which is hereby incorporated by reference in its entirety.

In one embodiment, sensor 212 includes a heart rate variability detector. Patients suffering acute decompensated heart failure exhibit abnormally low heart rate variability. An example of detecting the heart rate variability is discussed in U.S. Patent No. 5,603,331, "DATA LOGGING SYSTEM FOR  
5 IMPLANTABLE CARDIAC DEVICE," assigned to Cardiac Pacemakers, Inc., which is incorporated herein by reference in their entirety. Event detector 213 detects an abnormal condition when the heart rate variability falls below a predetermined threshold level.

In one embodiment, sensor 212 includes a renal function sensor. Acute  
10 decompensated heart failure results in peripheral edema primarily because of fluid retention of the kidneys that follows the reduction in cardiac output. The fluid retention is associated with reduced renal output, decreased glomerular filtration, and formation of angiotensin. Thus, in one specific embodiment, a renal function sensor includes a renal output sensor to sense a signal indicative  
15 of the renal output. Event detector 213 detects an abnormal condition when the sensed renal output falls below a predetermined threshold. In another specific embodiment, a renal function sensor includes a filtration rate sensor to sense a signal indicative of the glomerular filtration rate. Event detector 213 detects an abnormal condition when the sensed glomerular filtration rate falls below a  
20 predetermined threshold. In yet another specific embodiment, a renal function sensor includes a chemical sensor to sense a signal indicative of angiotensin II levels. Event detector 213 detects the abnormal condition when the sensed angiotensin II levels exceed a predetermined threshold level.

In one embodiment, sensor 212 includes an acoustic sensor such as a  
25 heart sound sensor and/or a respiratory sound sensor. Arrhythmias and/or heart failure cause abnormal cardiac and pulmonary activity patterns and hence, deviation of heart sounds and respiratory sounds from their normal ranges of pattern and/or amplitude. Event detector 213 detects an abnormal condition when the heart sound or respiratory sound is out of its normal range. For  
30 example, detection of abnormal third heard sound (S3) amplitude is known to indicate heart failure. In one specific embodiment, event detector 213 detects an abnormal condition when the S3 amplitude exceeds a predetermined threshold level.

In one embodiment, sensor 212 includes a remodeling sensor to sense a

signal indicative a degree of myocardial remodeling. In one specific embodiment, the remodeling sensor includes two or more piezoelectric crystals incorporated in one or more leads of lead system 108 to sense a size of an injured myocardial region such as an infarct region. The size of the injured myocardial region is estimated based on spatial information sensed by the crystals and averaged over a predetermined period of time. In one embodiment, a substantial degree of change in the size of the injured region indicates a need to start, stop, or adjust the combined electrical and agent therapies. In another specific embodiment, sensor 212 includes a hypertrophic sensor to sense a signal indicative of a degree of myocardial hypertrophy, which indicates the progress of the remodeling process. In another specific embodiment, sensor 212 includes a chemical sensor to sense the change in expression or concentration of Endothelin-1 (ET-1), BNP, or p38MAPK, which are known to change during hypertrophy response. In a further embodiment, sensor processing circuit 215 includes an event detector to detect an abnormal condition when the degree of myocardial remodeling exceeds a predetermined threshold. The degree of myocardial remodeling is represented by one or more of the degree of change in the size of the injured region, the degree of myocardial hypertrophy, and the degree of the change in expression or concentration of Endothelin-1 (ET-1), BNP, or p38MAPK.

Embodiments of sensor 212 and event detector 213 are discussed in this document by way of example, but not by way of limitation. Other methods and sensors for directly or indirectly detecting an abnormal condition treatable by the gene/protein delivery may be employed with gene/protein delivery system 100.

Implantable CRM device 110 includes a hermetically sealed metal can to house at least portion of the electronics of the device. In one embodiment, sensor 212 resides within the metal can. In another embodiment, sensor 212 is outside of the metal can. In one embodiment, sensor 212 is incorporated into a lead system 108.

FIG. 3 is a block diagram showing a further embodiment of the circuit of portions of system 100 including implantable CRM device 110, lead system 108, gene/protein delivery device 130, and external system 155. Implantable CRM device 110 as shown in FIG. 3 includes pacing and defibrillation capabilities. In addition to gene/protein delivery, examples of therapies delivered by implantable

CRM device 110 include, but are not limited to, bradyarrhythmia pacing, anti-tachyarrhythmia pacing, atrial and/or ventricular cardioversion/defibrillation, CRT, RCT, and drug delivery. However, the pacing and defibrillation capabilities are not necessary for system 100 to perform gene/protein delivery, and hence, may be excluded from implantable CRM device 110. In other words, implantable CRM device 110 can be an implantable pacemaker and/or defibrillator with additional functions including control of gene/protein delivery, or it can be a dedicated implantable gene/protein delivery processor or controller.

10 In one embodiment, implantable CRM device 110 includes sensor 212, event detector 213, implant controller 214, pacing circuit 320, defibrillation circuit 324, and implant telemetry module 316. Pacing circuit 320 delivers pacing pulses to one or more cardiac regions as controlled by implant controller 214. Defibrillation circuit 324 delivers cardioversion or defibrillation shocks to one or more cardiac regions as controlled by implant controller 214. Sensor 212 senses a physiological signal indicative of a condition treatable with gene/protein delivery, and event detector 213 detects that condition, as discussed above with reference to FIG. 2. In one specific embodiment, in which implantable CRM device 110 provides for CRT and RCT pacing as well as defibrillation, implant controller 214 includes gene/protein delivery control module 215, CRT control module 321, RCT control module 322, defibrillation control module 323, and command receiver 326. Gene/protein delivery control module 215 generates a gene/protein delivery control signal in response to a condition detected by event detector 213 or a gene/protein delivery command received by command receiver 326. Command receiver 326 receives a gene/protein delivery command from external system 155 via telemetry link 140. CRT control module 321 controls the delivery of pacing pulses from pacing circuit 320 by executing a CRT algorithm. RCT control module 322 controls the delivery of pacing pulses from pacing circuit 320 by executing a RCT algorithm. Defibrillation control module 323 controls the delivery of cardioversion/defibrillation shocks from defibrillation circuit 324 when a tachyarrhythmic condition is detected. In one embodiment, defibrillation control module 323 includes an atrial defibrillation control module to control the delivery of cardioversion/defibrillation shocks to one or more of the atria. In one embodiment, defibrillation control module 323



includes a ventricular defibrillation control module to control the delivery of cardioversion/defibrillation shocks to one or more of the ventricles.

Lead system 108 includes one or more gene/protein delivery control leads, referenced as lead system 108A, and pacing leads, defibrillation leads, 5 pacing-defibrillation leads, or any combination of such leads, referenced as lead system 108B. Lead system 108A allows gene/protein delivery control module 215 to control implantable gene/protein delivery device 130. Lead system 108B allows sensing of electrical signals from various regions of heart 101 and/or delivery of pacing pulses and/or defibrillation shocks to various regions of heart 10 101. The various regions of heart 101 includes regions within or about the right atrium (RA), left atrium (LA), right ventricle (RV), and left ventricle (LV). In one embodiment, lead system 108B includes one or more transvenous leads each having at least one sensing-pacing or defibrillation electrode disposed within heart 101. In one embodiment, lead system 108B includes one or more 15 epicardial leads each having at least one sensing-pacing or defibrillation electrode disposed on heart 101. In one embodiment, lead system 108B includes at least one atrial defibrillation electrode disposed in or about one or both of the atria to allow atrial defibrillation. In one embodiment, lead system 108B includes at least one ventricular defibrillation electrode disposed in or about one 20 or both of the ventricles to allow ventricular defibrillation. In one embodiment, sensor 212 includes at least portions of lead system 108A or 108B. In another embodiment, sensor 212 is incorporated into lead system 108A or 108B.

External system 155 includes external telemetry module 352, external user input device 354, presentation device 356, and external controller 358. 25 These system components distribute in one or more of external device 150, network 160, or remote device 170, depending on design and medical considerations. User input device 354 receives commands and/or parameters from the user and/or the patient to control deliveries of therapy, including gene/protein delivery. Presentation device 356 displays or otherwise presents 30 signals acquired and/or conditions detected by implantable CRM device 110. External controller 358 controls the operation of external system 155. In one embodiment, external controller 358 further provides automatic control of operations of an implantable CRM device 110. In one embodiment, user input device 352 receives the gene/protein delivery command entered by the user

based on observations of the signals and/or conditions presented by presentation device 356. In another embodiment, user input device 352 receives the gene/protein delivery command entered by a patient when the patient physically senses a symptom indicative of an immediate need for the gene/protein therapy, or entered by a person near the patient who observes a symptom indicative of the immediate need for the gene/protein therapy. In a further embodiment, external controller 358 automatically analyzes the signals acquired and/or abnormal conditions detected by implantable CRM device 110, and generates a gene/protein delivery command when deemed necessary as a result of the analysis.

Telemetry link 140 is a wireless bidirectional data transmission link supported by implant telemetry module 316 and external telemetry module 352. In one embodiment, telemetry link 140 is an inductive couple formed when two coils – one connected to implant telemetry module 316 and the other connected to external telemetry module 352 – are placed near each other. In another embodiment, telemetry link 140 is a far-field radio-frequency telemetry link allowing implantable CRM device 110 and external system 155 to communicate over a telemetry range that is at least ten feet.

#### Implantable Iontophoresis Gene/Protein Delivery Device

The system of the invention includes an implantable iontophoresis gene and/or protein delivery device coupled to an implantable pulse generator. The implantable iontophoresis gene/protein delivery device includes a polymer matrix and isolated nucleic acid which encodes at least one gene product; isolated nucleic acid which binds at least one selected mRNA, and/or isolated protein, or any combination thereof. The isolated nucleic acid or protein is selected at the discretion of the practitioner on the basis of a correlation of the sequence with a particular condition, e.g., a correlation with beneficially altering the condition. Examples of selected genes and proteins are provided below.

#### Exemplary Gene(s) and/or Protein(s)

For iontophoresis techniques to be used, the molecule embedded in or applied to the polymer matrix should have specific characteristics. Ideally, the molecule should have an ionic nature or have other ionic molecules bound to the molecule to promote the iontophoretic movement or transport of that molecule from the polymer matrix. Gene(s) and/or protein(s) suitable for the implantable

iontophoresis gene/protein delivery device of the invention include those useful to treat, inhibit (reduce) or eliminate one or more symptoms associated with a particular condition. In one embodiment, where an increase in a particular gene product is indicated to treat, inhibit or eliminate one or more symptoms

5 associated with a particular condition, at least a portion of an open reading frame encoding the gene product in sense orientation is operably linked to transcriptional control elements, for instance, a heterologous transcriptional control element, optionally including a tissue-specific control element, to form an expression cassette. For example, in one embodiment, the expression cassette

10 includes an open reading frame for connexin 43 operably linked to a cardiac-specific promoter. In another embodiment, the expression cassette includes an open reading frame for a mammalian ion channel protein which is operably linked to a viral promoter.

In another embodiment, where a decrease in a particular gene product is

15 indicated to treat, inhibit or eliminate one or more symptoms associated with a particular condition, at least a portion of an open reading frame for that gene product in antisense orientation is operably linked to transcriptional control elements, for instance, a heterologous transcriptional control element, optionally including a tissue-specific control element, to form an expression cassette.

20 Alternatively, antisense nucleic acid is not present in an expression cassette, i.e., not operably linked to a transcriptional control element, e.g., a promoter.

For purposes of the present invention, control elements, such as muscle-specific and inducible promoters, enhancers and the like, will be of particular use. Such control elements include, but are not limited to, those derived from the

25 actin and myosin gene families, such as from the myoD gene family (Weintraub et al., Science, 251, 761 (1991)); the myocyte-specific enhancer binding factor MEF-2 (Cserjesi and Olson, Mol. Cell Biol., 11, 4854 (1991)); control elements derived from the human skeletal actin gene (Muscat et al., Mol. Cell Bio., 7, 4089 (1987)) and the cardiac actin gene; muscle creatine kinase sequence

30 elements (Johnson et al., Mol. Cell Biol., 9, 3393 (1989)) and the murine creatine kinase enhancer (mCK) element; control elements derived from the skeletal fast-twitch troponin C gene, the slow-twitch cardiac troponin C gene and the slow-twitch troponin I gene; hypoxia-inducible nuclear factors (Semenza et al., Proc. Natl. Acad. Sci. USA, 88, 5680 (1991); Semenza et al., J. Biol. Chem.,

269, 23757); steroid-inducible elements and promoters, such as the glucocorticoid response element (GRE) (Mader and White, Proc. Natl. Acad. Sci. USA, 90, 5603 (1993)); the fusion consensus element for RU486 induction; and elements that provide for tetracycline regulated gene expression (Dhawan et al., Somat. Cell. Mol. Genet., 21, 233 (1995); Shockett et al., Proc. Natl. Acad. Sci. USA, 92, 6522 (1995)).

In one embodiment, cardiac cell restricted promoters will be of particular use. Cardiac specific promoters include, but are not limited to, promoters from the following genes: a  $\alpha$ -myosin heavy chain gene, e.g., a ventricular  $\alpha$ -myosin heavy chain gene,  $\beta$ -myosin heavy chain gene, e.g., a ventricular  $\beta$ -myosin heavy chain gene, myosin light chain 2v gene, e.g., a ventricular myosin light chain 2 gene, myosin light chain 2a gene, e.g., a ventricular myosin light chain 2 gene, cardiomyocyte-restricted cardiac ankyrin repeat protein (CARP) gene, cardiac  $\alpha$ -actin gene, cardiac m2 muscarinic acetylcholine gene, ANP gene, BNP gene, cardiac troponin C gene, cardiac troponin I gene, cardiac troponin T gene, cardiac sarcoplasmic reticulum Ca-ATPase gene, skeletal  $\alpha$ -actin gene, as well as an artificial cardiac cell-specific promoter.

Further, chamber-specific promoters or enhancers may also be employed, e.g., for atrial-specific expression, the quail slow myosin chain type 3 (MyHC3) or ANP promoter, or the cGATA-6 enhancer, may be employed. For ventricle-specific expression, the iroquois homeobox gene may be employed. Examples of ventricular myocyte-specific promoters include a ventricular myosin light chain 2 promoter and a ventricular myosin heavy chain promoter.

Other sources for promoters and/or enhancers are promoters and enhancers from the Csx/NKX 2.5 gene, titin gene,  $\alpha$ -actinin gene, myomesin gene, M protein gene, cardiac troponin T gene, RyR2 gene, Cx40 gene, and Cx43 gene, as well as genes which bind Mef2, dHAND, GATA, CarG, E-box, Csx/NKX 2.5, or TGF-beta, or a combination thereof.

Tissue-specific enhancers may also be employed. For instance, a preferred atrial-specific enhancer is the cGATA-6 enhancer. In other embodiments, the enhancer is not tissue-specific.

In yet other embodiments, the promoter is a non-muscle, non-cardiac tissue-specific promoter.

Nevertheless, other promoters and/or enhancers which are not specific for certain tissue or cells, e.g., a viral promoter such a one from an adenovirus, adeno-associated virus, retrovirus, herpesvirus or lentivirus, may be employed in the expression cassettes and methods of the invention. A preferred heterologous  
 5 constitutive promoter is a CMV promoter which also includes an enhancer.

Preferably, the gene is also operably linked to a polyadenylation signal.

In one embodiment, the condition is a cardiovascular condition including, but not limited to, atrial fibrillation, ischemia, ventricular tachycardia, bradycardia, hyperplasia, or heart failure. Genes and corresponding proteins  
 10 useful in the system and methods for cardiovascular conditions include, but are not limited to, a connexin gene, an atrial-specific ion channel protein gene, e.g., a gene product associated with  $I_f$ , a non-specific ion channel protein gene, e.g., a gene product associated with  $I_K$ , a gene product which regulates gap junctions, a gene product which alters conduction in the myocardium, a gene product which  
 15 encodes a regulatory protein which blocks ion channel protein synthesis or activity, a gene product which upregulates ion channel protein synthesis or activity, a gene product which downregulates at least one ion channel protein, or a gene product which alters ion channel kinetics or voltage.

Thus, useful genes and corresponding proteins include genes for ion  
 20 channels, including  $K^+$ ,  $Na^+$ ,  $Ca^+$  and voltage activated channels, and preferably ion channels expressed in cardiac tissue, including, but not limited to, HCN (for  $I_f$ ), Kir 2.1 (for  $I_{K1}$ ), Kir 3.1/3.4 (for  $I_{KACh}$ ), ERG ( $\alpha$  subunit for  $I_{Kr}$ ), MiRP1 (modulates  $I_{Kr}$ ,  $I_f$  and  $I_{to}$ ), KvLQT1 ( $\alpha$  subunit for  $I_{Ks}$ ), MinK ( $\beta$  subunit for  $I_{Ks}$ ), Kv4.2/4.3 ( $\alpha$  subunit for  $I_{to}$ ), Kv1.4 ( $\alpha$  subunit for  $I_{to}$ ), KChIP2 ( $\beta$  subunit for  
 25  $I_{to}$ ), Kv1.5 (for  $I_{Kur}$ ), Cav1.2 ( $I_{CaL}$ ), Cav1.3 ( $I_{CaV}$ ), Cav3.1 ( $I_{Ca}$ ), Nav1.5 ( $I_{Na}$ ), and connexin 40, connexin 43, and connexin 45 ( $I_{GJ}$ ) (see Schram et al., Circ. Res., 90:939 (2002)), the disclosure of which is specifically incorporated by reference herein), NCX, e.g., NCX1, NCX2 and NCX3, Kir 6.1, Kir6.2, Kv1.7, Kv4.2, Kv4.3, Kv4.1 as well as Na ion channels such as SCN5A. Other genes useful in  
 30 the system and methods of the invention include, but are not limited to, Galphai2 subunit, neurotensin, calmodulin, calmodulin-dependent kinase, ATF3, calcitonin gene related peptide (CGRP), NOS, e.g., nNOS, iNOS and eNOS, Nkx 2.5, dystrophin, tafazzin, cardiac actin, desmin, lamin A/C, delta sarcoglycan, cardiac  $\beta$  myosin heavy chain, and cardiac tropin C,  $\beta$ -adrenergic

receptors, inhibitory guanine nucleotide protein, GPCR kinases, VEGF, placental growth factor, ACE(2), Cox2, for calcium regulation, calmodulin, CaMKII, RyR (ryanodine receptor), SERCA, calcium ATPase (CSR), phospholamban (PLB), calcineurin, and FK506 binding protein, Lbx1, AT1A receptor, AT2, p27(KIP1), calcineurin, angiotensin IT, and HSP. To prevent, inhibit or treat heart failure, the following genes/proteins may be useful: calmodulin, CaMKII, RyR (ryanodine receptor), SERCA, calcium ATPase (CSR), phospholamban (PLB), calcineurin, and FK506 binding protein. To prevent, inhibit or treat hyperplasia, the following genes/proteins may be useful: Lbx1, AT1A receptor, AT2, p27(KIP1), calcineurin, angiotensin II, and HSP. To prevent, inhibit or treat ischemia, the following genes/proteins may be useful: CGRP, ATF3, placental growth factor and Cox2. To increase contractility, the following genes/proteins may be useful: neurotensin and ACE(2).

#### Polymer Matrix

The polymer matrix may be formed of any physiologically compatible material which generally retains isolated nucleic acid and/or protein (which are charged molecules) or optionally other agents including other therapeutic agents under physiological conditions for a sustained period of time, e.g., for months or years, in the absence of an electrical field. The polymer matrix extrudes (releases) isolated nucleic acid or protein from the implantable iontophoresis gene or protein delivery device in response to an electric field created by an electrical signal. The electric signal is generated in response to the detection of a physiological signal associated with a condition, e.g., a cardiovascular condition.

The isolated nucleic acid and/or protein or optional other agent(s) may be introduced to a solution of monomers prior to polymerization or to the polymer matrix, e.g., dissolved in a solvent (e.g., water, propylene, glycol, etc.) and the resulting solution can be incorporated into the polymer matrix material. Once the isolated nucleic acid and/or protein is embedded in or applied to a polymer matrix, the resulting gene/protein delivery device can be coupled to an implantable pulse generator. Alternatively, the polymer matrix may be first coupled to the implantable pulse generator and then the isolated nucleic acid and/or protein embedded in or applied thereto, either passively or actively (through, for example, such methods as iontophoresis). Upon delivery of an electric field, the isolated nucleic acid and/or protein or optional other agent(s) is

released from the matrix at a rate which is greater than the rate of release in the absence of the electric field. In particular, the isolated nucleic acid and/or protein in an implanted iontophoresis gene/protein delivery device is released to adjacent cells or tissue or the vessel lumen in response to an electric field  
5 generated by the implantable pulse generator, which release is in an amount proportional to the applied electric field. Once the electric signal is stopped, the genes and/or proteins are no longer released or released at a rate which is significantly reduced relative to the rate of release in the presence of the electric field. Thus, the delivery of the gene(s) and/or protein(s) is transient (temporary)  
10 and may be spatially controlled by the direction of the electric field and placement of the device.

The matrix materials will preferably be physiologically inert and capable of retaining the charged molecule to be delivered. Matrix materials which may be used include: polyacetic or polyglycolic acid and derivatives thereof,  
15 polyorthoesters, polyesters, polyurethanes, polyamino acids such as polylysine, lactic/glycolic acid copolymers, polyanhydrides and ion exchange resins such as sulfonated polytetrafluorethylene, or combinations thereof.

Additionally, it is possible to construct the matrices from natural proteins or materials which are crosslinked using a crosslinking agent such as 1-ethyl-3-  
20 (3-dimethylamino-propyl)carbodiimide hydrochloride. Such natural materials are those such as albumin, collagen, fibrin, gelatin, keratin, potato starch hydrolyzed for use in electrophoresis, and agar-agar (agarose). Synthetic electrophoretic matrices are also suitable including polyacrylamide, acrylamide/bis-acrylamide mixtures, cellulose acetate, glyoxyl agarose, and  
25 Sephadex™ (Pharmacia Fine Chemicals, Inc.) suitable for use in isoelectrofocusing. It is also possible to use combinations of such matrices, such as the combination of polyacrylamide and agarose, in order to fabricate the cathodic matrix of the invention.

In one embodiment, the polymer matrix may include liposomes, a  
30 hydrogel, cyclodextrins, biodegradable nanocapsules or microspheres. Thus, polymer matrix includes synthetic polymers in the form of hydrogels or other porous materials, e.g., nucleic acid and/or protein permeable configurations or morphologies, such as polyvinyl alcohol, polyvinylpyrrolidone and polyacrylamide, polyethylene oxide, poly(2-hydroxyethyl methacrylate); natural

polymers such as gums and starches; synthetic elastomers such as silicone rubber, polyurethane rubber; and natural rubbers, and include poly[ $\alpha$ -(4-aminobutyl)]-L-glycolic acid, polyethylene oxide (Roy et al., Mol. Ther., 7:401 (2003)), poly orthoesters (Heller et al., Adv. Drug Delivery Rev., 54:1015 (2002)), silk-elastin-like polymers (Megeld et al., Pharma. Res., 19:954 (2002)), alginate (Wee et al., Adv. Drug Deliv. Rev., 31:267 (1998)), EVAc (poly(ethylene-co-vinyl acetate), microspheres such as poly (D, L-lactide-co-glycolide) copolymer and poly (L-lactide), poly(N-isopropylacrylamide)-b-poly(D,L-lactide), a soy matrix such as one cross-linked with glyoxal and reinforced with a bioactive filler, e.g., hydroxylapatite, poly(epsilon-caprolactone)-poly(ethylene glycol) copolymers, poly(acryloyl hydroxyethyl) starch, polylysine-polyethylene glycol, an agarose hydrogel, or a lipid microtubule-hydrogel.

In one embodiment, the nucleic acid or protein is embedded in or applied to a polymer matrix, e.g., a nonionic or ionic biodegradable or nonbiodegradable matrix, including but not limited to hydrogels of poloxamers, polyacrylamide, poly(2-hydroxyethyl methacrylate), carboxyvinyl-polymers (e.g., Carbopol 934, Goodrich Chemical Co.), cellulose derivatives, e.g., methylcellulose and hydroxypropyl cellulose, polyvinyl pyrrolidone or polyvinyl alcohols.

In another embodiment, the biocompatible polymeric materials are synthetic, nonbiodegradable polymers such as polyurethanes, polydimethylsiloxanes (silicone rubbers), ethylene vinyl acetate copolymer (EVA), poly methylmethacrylate, polyamides, polycarbonates, polyesters, polyethylene, polypropylenes, polystyrenes, polyvinyl chloride, polyvinyl alcohols, polytetrafluoroethylene, or cellulose derivatives such as cellulose acetate.

In alternative embodiments, the biocompatible polymeric material is a biodegradable polymeric such as collagen, fibrin, polylactic-polyglycolic acid, or a polyanhydride. Other examples include, without limitation, any biocompatible polymer, whether hydrophilic, hydrophobic, or amphiphilic, such as ethylene vinyl acetate copolymer (EVA), polymethyl methacrylate, polyamides, polycarbonates, polyesters, polyethylene, polypropylenes, polystyrenes, polyvinyl chloride, polytetrafluoroethylene, or cellulose derivatives such as cellulose acetate. In an alternative embodiment, a biologically derived polymer,



such as protein collagen, fibrin, polylactic-polyglycolic acid, or a polyanhydride, is a suitable polymeric matrix material.

The above examples are provided for reference only, and the range of suitable polymer matrix materials should not be construed as limited to those materials listed above. The polymer matrix material can be hydrophilic, hydrophobic, or amphiphilic, provided it meets the physical characteristics described above. See also U.S. Patent No. 5,087,243 and Avitall et al., Circ., 85:1582 (1992). The polymer matrix preferably stabilizes the gene(s) and/or protein(s) and other optional agents in the polymer matrix.

10 A polymer matrix may also be present in a selectively semipermeable membrane such as a dialysis membrane, nylon or polysulfoxy. In one embodiment, the semipermeable membrane is not biodegradable.

In another embodiment, it is possible to fabricate multiple-layered cathodic reservoirs. Such multiple-layered reservoirs will find usefulness in certain embodiments in which there is desire to have additional control over the rate at which the charged substance is electrophoresed from the matrix. Thus, where it is desired to prevent diffusion of an anionic molecule chiefly occurring at the surface of the reservoir in contact with the target tissue, a relatively strongly-cationic matrix material may be used to cap a relatively weakly-cationic matrix material. This multiple-layered matrix embodiment may be preferentially utilized where a relatively more toxic, but more efficacious molecule is to be initially utilized to lower the immediate post-implantation stimulation threshold. However, and particularly in patients with a history of cardiac arrhythmia or fibrillation, it may be useful to charge a second layer with an anti-arrhythmic or antifibrillation drug, which drug is delivered only upon the necessary physiological demand. Similar combinations may be made in patients combinations may be made in patients with histories of cardiac arrest, where thyroid hormone therapy may be desired.

When a multiple-layered matrix is used, it is also possible to sequentially deliver a gradually declining (or increasing, or cyclical) concentration of the molecule. In such an embodiment, the distal most matrix reservoir may contain a first lower dose of the molecule, followed by a next most distal matrix with a concentration of the molecule higher than the first, and so on.

#### Methods of Using the Implantable Systems

The implantable systems of the invention which include an implantable pulse generator and an implantable iontophoresis gene/protein delivery device may coupled to other implantable devices, e.g., a stent, shunt, indwelling catheter, lead or epicardial patch, or minimally invasive devices, e.g., a

5 transdermal patch. Such combined devices may be introduced to vessels, e.g., a vein, or other body lumens. In one embodiment, an implantable system of the invention is introduced to a mammalian heart, e.g., to one or both atria. For example, a mammal having or at risk of having atrial fibrillation is provided with a system which includes an implantable pulse generator and an implantable

10 iontophoresis gene/protein delivery device, which system may be coupled to the atria of the heart. Upon sensing a physiological signal indicative of atrial fibrillation, the implant controller in the implantable pulse generator produces an electric signal which results in an electric field. The isolated nucleic acid and/or protein in the implantable iontophoresis gene/protein delivery device is released

15 to adjacent cells or tissue or the vessel lumen in response to the electric field in an amount proportional to the applied electric field. The dosage ranges for the gene(s)/protein(s) are those large enough to produce the desired effect in which the symptoms of the condition are ameliorated. The dosage should not be so large as to cause adverse side effects. Generally, nucleic acid dosages include

20 from about 0.001 to about 500  $\mu\text{g}$ , e.g., from about 0.01  $\mu\text{g}$  to about 100  $\mu\text{g}$ , and for protein dosages, from about 1 ng to about 10 mg, e.g., from about 1  $\mu\text{g}$  to about 1 mg. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient and can be determined by one of skill in the art. Once the electric signal is stopped, the genes and/or proteins are no longer

25 released or released at a rate which is significantly reduced relative to the rate of release in the presence of the electric field. Thus, the delivery of the gene(s) and/or protein(s) is transient (temporary) and may be spatially controlled by the direction of the electric field and placement of the device.

In one embodiment, where the heart tissue around the entire periphery of

30 the heart is to receive the nucleic acid/protein, it is administered to the pericardial sac. In another embodiment, nucleic acid/protein which is present in a hydrogel can be percutaneously applied to the surface of the heart applied to the heart muscle. However, it will be appreciated that the strength and/or

distribution of the electric field can be controlled such that a larger or smaller area of the heart tissue can be treated.

5       The frequency range for iontophoresis begins at 0 Hz (dc) and increases to a maximum of about 20 MHz, with the preferred range lying between 2-15 kHz. In one embodiment, the electric field is an ac field with a dc offset. It will be understood that the frequency can be varied within these ranges to maximize the rate of iontophoretic transfer for a given drug used in the catheters of the present invention. Selecting a frequency range that is significantly higher than the intrinsic heart rate reduces the risk of inducing an arrhythmia.

10       The apparatus and system of the invention may be employed with other therapies, e.g., electrical therapies such as pacing, CRT, RCT and defibrillation, drug therapies, and/or non-localized gene therapy, e.g., which results in integrated genes. Other therapies may include but are not limited to treatment with growth factors and/or angiogenic factors, or drugs such as beta-adrenergic  
15       blockers, e.g., propranolol, calcium channel blockers, e.g., verapamil, angiotensin converting enzyme inhibitors, e.g., captopril, angiotension II receptor blockers, e.g., losartan, alpha adrenergic blockers, e.g., doxazosin, or hypotensive agents, e.g., reserpine, antilipemic agents, e.g., simvastatin, vasodilating agents, e.g., amyl nitrite, carvedilol, adenosine, digoxin, ibutilide,  
20       lidocaine, neseritide and the like.

      All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification, this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled  
25       in the art that the invention is susceptible to additional embodiments and that certain of the details herein may be varied considerably without departing from the basic principles of the invention.

**WHAT IS CLAIMED:**

1. A system, comprising:
  - an implantable pulse generator including:
    - a sensor to sense a physiological signal indicative of a predetermined cardiac condition;
    - an event detector, coupled to the sensor, to detect the predetermined cardiac condition from the physiological signal; and
    - an implant controller coupled to the event detector, the implant controller including a gene or protein delivery control module adapted to produce an electrical signal to control gene or protein delivery in response to the predetermined cardiac condition; and
  - an implantable gene or protein delivery device coupled to the implantable pulse generator, the implantable gene or protein delivery device including a polymer matrix and isolated nucleic acid which encodes at least one gene product or binds at least one selected mRNA, or isolated protein, which nucleic acid or protein is released from the implantable gene or protein delivery device in response to an electric field created by the electrical signal.
2. The system of claim 1, wherein the sensor comprises an electrogram sensing circuit, and the event detector comprises an arrhythmia detector.
3. The system of claim 2, wherein the arrhythmia detector comprises one or more of a bradycardia detector, a tachycardia detector, and a fibrillation detector.
4. The system of claim 3, wherein the fibrillation detector comprises an atrial fibrillation detector.
5. The system of any one of claims 3 to 4, wherein the fibrillation detector comprises a ventricular fibrillation detector.

6. The system of claim 1, wherein the sensor comprises a sensor sensing an physiological signal indicative of ischemia, and the event detector comprises an ischemia detector.
7. The system of claim 1, wherein the sensor comprises a metabolic sensor adapted to sense a signal indicative of a cardiac metabolic level.
8. The system of claim 7, wherein the sensor comprises at least one of a pH sensor, an oxygen pressure (PO<sub>2</sub>) sensor, a carbon dioxide pressure (PCO<sub>2</sub>) sensor, a glucose sensor, a creatine sensor, a C-creative protein sensor, a creatine kinase sensor, and a creatine kinase-MB sensor.
9. The system of claim 1, wherein the sensor comprises an impedance sensor to sense tissue impedance.
10. The system of claim 9, wherein the impedance sensor comprises a pulmonary impedance sensor.
11. The system of claim 10, wherein the impedance sensor comprises a respiratory sensor.
12. The system of claim 1, wherein the sensor comprises a pressure sensor to sense a pressure in a cardiovascular system.
13. The system of claim 12, wherein the pressure sensor comprises at least one of a left atrial pressure sensor, a left ventricular pressure sensor, an artery pressure sensor, and a pulmonary arterial pressure sensor.
14. The system of any one of claims 12 to 13, wherein the event detector comprises a systolic dysfunction detector.

15. The system of any one of claims 12 to 14, wherein the event detector comprises a diastolic dysfunction detector.
16. The system of claim 1, wherein the sensor comprises a stroke volume sensor.
17. The system of claim 1, wherein the sensor comprises a neural activity sensor.
18. The system of claim 17, wherein the neural activity sensor comprises a neurohormone sensor to sense a neurohormone level.
19. The system of claim 17, wherein the neural activity sensor comprises a action potential recorder to sense neural electrical activities.
20. The system of claim 1, wherein the sensor comprises a heart rate variability detector.
21. The system of claim 1, wherein the sensor comprises a renal function sensor.
22. The system of claim 21, wherein the renal function sensor comprises at least one of a renal output sensor, a filtration rate sensor, and an angiotensin II level sensor.
23. The system of claim 1, wherein the sensor comprises an acoustic sensor adapted to sense at least one of heart sounds and respiratory sounds.
24. The system of claim 23, wherein the event detector to detect the predetermined cardiac condition when third hear sound (S3) amplitude exceeds a predetermined threshold.
25. The system of claim 1, wherein the sensor comprises a remodeling sensor to sense a signal indicative a degree of myocardial remodeling.

26. The system of claim 25, wherein the remodeling sensor comprises two or more piezoelectric crystals to sense a size of an injured myocardial region.
27. The system of any one of claims 1 to 26, wherein the implantable gene or protein delivery device comprises an epicardial patch including the polymer matrix.
28. The system of any one of claims 1 to 27, wherein the nucleic acid or protein is released at a rate controlled by the strength of the electric field.
29. The system of any one of claims 1 to 28, wherein the implantable pulse generator further comprises a pacing circuit coupled to the implant controller, and wherein the implant controller includes a pacing control module adapted to control a delivery of pacing pulses in conjunction with the release of the nucleic acid or protein.
30. The system of claim 29, further comprising at least one atrial pacing lead coupled to the pacing circuit to deliver atrial pacing pulses.
31. The system of any one of claims 29 to 30, further comprising at least one ventricular pacing lead coupled to the pacing circuit to deliver ventricular pacing pulses.
32. The system of any one of claims 29 to 31, wherein the implantable pulse generator further comprises a cardiac resynchronization therapy (CRT) circuit coupled to the implant controller, and wherein the implant controller includes a CRT control module adapted to control a delivery of CRT in conjunction with the release of the nucleic acid or protein.
33. The system of any one of claims 29 to 32, wherein the implantable pulse generator further comprises a remodeling control therapy (RCT) circuit coupled to the implant controller, and wherein the implant controller includes an RCT therapy

control module adapted to control a delivery of RCT therapy in conjunction with the release of the nucleic acid or protein.

34. The system of any one of claims 1 to 33, wherein the implantable pulse generator further comprises defibrillation circuit coupled to the implant controller, and wherein the implant controller includes a defibrillation control module adapted to control a delivery of defibrillation shocks in conjunction the release of the nucleic acid or protein.

35. The system of claim 34, further comprising at least one atrial defibrillation lead coupled to the defibrillation circuit to deliver the defibrillation shocks to one or more atria, and wherein the defibrillation control module comprises an atrial defibrillation control module.

36. The system of any one of claims 34 to 35, further comprising at least one ventricular defibrillation lead coupled to the defibrillation circuit to deliver the defibrillation shocks to one or more ventricles, and wherein the defibrillation control module comprises a ventricular defibrillation control module.

37. The system of any one of claims 1 to 36, wherein the implantable pulse generator further comprises an implant telemetry module to receive an external command, and wherein the gene or protein delivery control module is adapted to produce the electrical signal to control gene or protein delivery in response to the predetermined cardiac condition and the external command.

38. The system of claim 37, further comprising an external system communicatively coupled to the implantable pulse generator, the external system including:

- a presentation device to present one or more of the sensed physiological signal and the detected predetermined cardiac condition;
- a user input device to receive the external command; and



an external telemetry module to transmit the external command to the implant telemetry module.

39. The system of claim 38, wherein the external system comprises a programmer.

40. The system of claim 38, wherein the external system comprises an advanced patient management system including:

an external device wirelessly coupled to the implantable pulse generator via telemetry;

a remote device to provide for access to the implantable pulse generator from a distant location; and

a network connecting the external device and the remote device.

41. The system of claim 40, wherein the external device comprises the user input.

42. The system of claim 40, wherein the remote device comprises the user input.

43. A method for operating an implantable gene or protein delivery system, comprising:

sensing a physiological signal indicative of a predetermined cardiac condition using an implantable sensor;

detecting the predetermined cardiac condition from the physiological signal;

and

producing an electrical signal in response to the predetermined cardiac condition, which electric signal is effective to create an electric field which is capable of releasing nucleic acid or protein from an implantable gene or protein delivery device including a polymer matrix and isolated nucleic acid which encodes at least one gene product or binds at least one selected mRNA or isolated protein.

44. The method of claim 43, wherein the nucleic acid or protein is released at a rate controlled by the strength of the electric field.

45. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing at least one electrogram, and detecting the predetermined cardiac condition comprises detecting an arrhythmia.

46. The method of claim 45, wherein detecting the predetermined cardiac condition comprises detecting an atrial fibrillation.

47. The method of claim 45, wherein detecting the predetermined cardiac condition comprises detecting a ventricular fibrillation.

48. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing an physiological signal indicative of ischemia, and detecting the predetermined cardiac condition comprises detecting an ischemia.

49. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a signal indicative of a cardiac metabolic level.

50. The method of claim 49, wherein sensing the signal indicative of the cardiac metabolic level comprises sensing at least one of a pH value, an oxygen pressure ( $\text{PO}_2$ ), a carbon dioxide pressure ( $\text{PCO}_2$ ), a glucose level, a creatine level, a C-creative protein level, a creatine kinase level, and a creatine kinase-MB level.

51. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing tissue impedance.

52. The method of claim 51, wherein sensing the tissue impedance comprises sensing pulmonary impedance.

53. The method of claim 51, wherein sensing the tissue impedance comprises sensing an impedance indicative of minute ventilation.

54. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a pressure in a cardiovascular system.

55. The method of claim 54, wherein sensing the pressure comprises sensing at least one of a left atrial pressure, a left ventricular pressure, an arterial pressure, and a pulmonary arterial pressure.

56. The method of claim 54, wherein detecting the predetermined cardiac condition comprises detecting a systolic dysfunction.

57. The method of claim 54, wherein detecting the predetermined cardiac condition comprises detecting a diastolic dysfunction.

58. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a stroke volume.

59. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a neural activity.

60. The method of claim 59, wherein sensing the neural activity comprises sensing a neurohormone level.

61. The method of claim 59, wherein sensing the neural activity comprises sensing neural electrical activities.

62. The method of claim 61, wherein sensing the physiological signal comprises detecting a heart rate variability.

63. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a renal function.

64. The method of claim 63, wherein sensing the renal function comprises sensing at least one of a renal output, a filtration rate, and an angiotensin II level.

65. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing at least one of heart sounds and respiratory sounds.

66. The method of claim 65, wherein detecting the predetermined cardiac condition comprises detecting a predetermined cardiac condition when third heart sound (S3) amplitude exceeds a predetermined threshold.

67. The method of any one of claims 43 to 44, wherein sensing the physiological signal comprises sensing a signal indicative of a degree of myocardial remodeling.

68. The method of claim 67, wherein sensing the signal indicative of the degree of myocardial remodeling comprises sensing a size of an injured myocardial region.

69. The method of any one of claims 43 to 68, further comprising delivering pacing pulses in conjunction with releasing the nucleic acid or protein.

70. The method of claim 69, further comprising delivering a cardiac resynchronization therapy (CRT) in conjunction with releasing the nucleic acid or protein.

71. The method of any one of claims 69 to 70, further comprising delivering a remodeling control therapy (RCT) in conjunction with releasing the nucleic acid or protein.

72. The method of any one of claims 43 to 71, further comprising delivering cardioversion/defibrillation shocks in conjunction with releasing the nucleic acid or protein.

73. The method of claim 72, wherein delivering the cardioversion/defibrillation shocks comprises delivering atrial defibrillation shocks.

74. The method of any one of claims 72 to 73, wherein delivering the cardioversion/defibrillation shocks comprises delivering ventricular defibrillation shocks.

75. The method of any one of claims 43 to 74, further comprising receiving an external command transmitted to an implantable device from an external system, and wherein producing the electrical signal comprises producing the electrical signal in response to the predetermined cardiac condition and the external command.

76. The method of claim 75, further comprising:  
transmitting one or more of the sensed physiological signal and a detection of the predetermined cardiac condition to the external system; and  
presenting the one or more of the sensed physiological signal and a detection of the predetermined cardiac condition through the external system.

77. The method of any one of claims 75 to 76, where receiving the external command comprises receiving the external command entered by a physician or other caregiver through the external system.

78. The method of any one of claims 75 to 76, where receiving the external command comprises receiving the external command entered by a patient through the external system.

79. A method to prepare an implantable gene or protein delivery device, comprising:  
introducing to a physiologically compatible polymer matrix isolated nucleic acid which encodes at least one gene product or binds at least one selected mRNA in a mammal or isolated protein, so as to yield an implantable gene or protein delivery

device, wherein the isolated nucleic acid or protein is selected so as to prevent, inhibit or treat a cardiac condition in a mammal, and wherein the rate of release of the nucleic acid or protein from the device is capable of being controlled by an electric signal.

80. A method to prepare an implantable gene or protein delivery device, comprising:

combining physiologically compatible monomers suitable to form a polymeric matrix and isolated nucleic acid which encodes at least one gene product or binds at least one selected mRNA in a mammal or isolated protein, so as to yield an implantable gene or protein delivery device, wherein the isolated nucleic acid or protein is selected so as to prevent, inhibit or treat a cardiac condition in a mammal, and wherein the rate of release of the nucleic acid or protein from the device is capable of being controlled by an electric signal.

81. The method of claim 79 or 80, wherein the nucleic acid or protein is selected to prevent, inhibit or treat atrial fibrillation, heart failure, ventricular fibrillation, ischemia, brachycardia or hyperplasia.

82. The method of claim 79 or 80, wherein the nucleic acid is present on one or more vectors.

83. The method of claim 82, wherein the one or more vectors are DNA vectors.

84. The method of claim 82, wherein the one or more vectors are viral vectors.

85. The method of claim 82, wherein the one or more vectors are plasmid vectors.

86. The method of any one of claims 79 to 81, wherein the nucleic acid includes morpholino modified nucleotides.

87. The method of any one of claims 79 to 86, wherein at least one gene product or protein is a connexin.
88. The method of any one of claims 79 to 86, wherein at least one gene product or protein is an atrial-specific ion channel protein.
89. The method of claim 88, wherein the gene product or protein is associated with  $I_f$ .
90. The method of any one of claims 79 to 86, wherein at least one gene product or protein is a non-specific ion channel protein.
91. The method of claim 90, wherein the gene product or protein is associated with  $I_{K1}$ ,  $I_{CaL}$ ,  $I_{to}$ ,  $I_{Kr}$ ,  $I_{Kur}$ ,  $I_{KATP}$ ,  $I_{Na}$  or  $I_{Na/Ca}$ .
92. The method of any one of claims 79 to 86, wherein at least one gene product or protein regulates gap junctions.
93. The method of any one of claims 79 to 86, wherein at least one gene product or protein alters conduction in the myocardium.
94. The method of any one of claims 43 or 79 to 86, wherein the protein is or the nucleic acid encodes a regulatory protein that blocks ion channels, upregulates ion channels, downregulates ion channels, or alters ion channel kinetics.
95. An implantable gene or protein delivery device prepared by the method of any one of claims 79 or 94.
96. A method to transiently deliver isolated nucleic acid which encodes at least one gene product or binds at least one selected mRNA, or isolated protein to a mammal, comprising:

providing a mammal comprising the system of any one of claims 1 to 42,  
wherein the mammal is at risk of the predetermined cardiac condition; and  
directing the production of the electric signal in response to detection of the  
condition so as to transiently deliver the nucleic acid or protein in an amount effective  
to inhibit or treat the condition or at least one symptom thereof.

97. The method of claim 96, wherein the system is in or on the heart of the mammal.

98. The method of claim 96, wherein the system is in or on a blood vessel of the mammal.

99. The method of any one of claims 96 to 98, wherein the condition is atrial fibrillation, heart failure, ventricular fibrillation, ischemia, brachycardia or hyperplasia.

100. The method of any one of claims 96 to 98, wherein the nucleic acid or protein is delivered to at least one of the atria.

101. The method of any one of claims 96 to 98, wherein the nucleic acid or protein is delivered to at least one of the ventricles.

102. The method of any one of claims 96 to 98, wherein the nucleic acid or protein delivered is a connexin nucleic acid or protein.

103. The method of any one of claims 96 to 98, wherein the nucleic acid or protein delivered is an atrial-specific ion channel gene or protein.

104. The method of claim 103, wherein the nucleic acid or protein delivered is associated with  $I_f$ .




105. The method of any one of claims 96 to 98, wherein the nucleic acid or protein delivered is a non-specific ion channel gene or protein.

106. The method of claim 105, wherein the nucleic acid or protein delivered is associated with  $I_{K1}$ ,  $I_{CaL}$ ,  $I_{to}$ ,  $I_{Kr}$ ,  $I_{Kur}$ ,  $I_{KATP}$ ,  $I_{Na}$  or  $I_{Na/Ca}$ .


107. The method of any one of claims 96 to 98, wherein the gene product or protein regulates gap junctions.

108. The method of any one of claims 96 to 98, wherein the gene product or protein alters conduction in the myocardium.

109. The method of any one of claims 96 to 98, wherein the protein is or the nucleic acid encodes a regulatory protein that blocks ion channels, upregulates ion channels, downregulates ion channels, or alters ion channel kinetics.



*Fig. 1*

*Fig. 2*




Fig. 3

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US2005/024914

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 A61N1/30		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61N A61M A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/74773 A (IMPULSE DYNAMICS N.V; DARVISH, NISSIM; SHEMER, ITSIK) 14 December 2000 (2000-12-14) page 4, line 12 - page 5, line 5 page 13, line 13 - page 14, line 2 page 16, line 5 - page 17, line 14 page 20, line 8 page 29, line 9 - line 25	1-42, 79-95
Y	WO 02/087681 A (MEDTRONIC, INC) 7 November 2002 (2002-11-07) page 3, line 3 - line 10 page 7, line 7 - line 24 page 11, line 6 - line 14 page 14, line 3 - line 14 page 33, line 7 - page 34, line 2 ----- -/--	1-42, 79-95
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search  3 November 2005		Date of mailing of the international search report  10/11/2005
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Petter, E

## INTERNATIONAL SEARCH REPORT

 International Application No  
 PCT/US2005/024914

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 087 243 A (AVITALL ET AL) 11 February 1992 (1992-02-11) abstract -----	1-42, 79-95
A	US 5 807 306 A (SHAPLAND ET AL) 15 September 1998 (1998-09-15) abstract -----	1,79,80, 95
X	WO 2004/026394 A (ADVANCED CARDIOVASCULAR SYSTEMS, INC) 1 April 2004 (2004-04-01) page 24, paragraph 3 - page 25, paragraph 2 page 16, paragraph 4 -----	79,80
A	WO 98/02150 A (MEDTRONIC, INC) 22 January 1998 (1998-01-22) abstract -----	1,79,80, 95
P,A	WO 2004/093969 A (MEDTRONIC, INC; LASKE, TIMOTHY, G; SIGG, DANIEL, C; SOYKAN, ORHAN) 4 November 2004 (2004-11-04) page 6, paragraph 4 - page 7, paragraph 1 page 8, paragraph 2 -----	1,79,80, 95
A	WO 00/62855 A (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 26 October 2000 (2000-10-26) abstract page 18, line 3 - line 18 -----	1,79,80, 95
A	WO 98/02040 A (MEDTRONIC, INC) 22 January 1998 (1998-01-22) abstract page 11, line 1 - page 13, line 12 -----	1,79,80, 95

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US2005/024914

### Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 43-78, 96-109  
because they relate to subject matter not required to be searched by this Authority, namely:  
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by surgery and therapy
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2005/024914

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0074773	A	14-12-2000	AU 4947100 A WO 0193951 A1 WO 0193950 A1	28-12-2000 13-12-2001 13-12-2001
WO 02087681	A	07-11-2002	EP 1385575 A1 EP 1385570 A2 WO 02087696 A1	04-02-2004 04-02-2004 07-11-2002
US 5087243	A	11-02-1992	AT 127028 T CA 2044791 A1 DE 69112533 D1 DE 69112533 T2 EP 0468636 A1 JP 2686383 B2 JP 5115562 A	15-09-1995 19-12-1991 05-10-1995 08-02-1996 29-01-1992 08-12-1997 14-05-1993
US 5807306	A	15-09-1998	NONE	
WO 2004026394	A	01-04-2004	AU 2003270723 A1 EP 1542762 A1 US 2004186546 A1	08-04-2004 22-06-2005 23-09-2004
WO 9802150	A	22-01-1998	AU 2458597 A CA 2260756 A1 EP 0957902 A1	09-02-1998 22-01-1998 24-11-1999
WO 2004093969	A	04-11-2004	NONE	
WO 0062855	A	26-10-2000	AU 4465600 A EP 1210144 A1	02-11-2000 05-06-2002
WO 9802040	A	22-01-1998	AU 2437397 A CA 2261022 A1 EP 0928135 A1 US 6567705 B1	09-02-1998 22-01-1998 14-07-1999 20-05-2003